

A Study of The Cohomology of The Orlik-Solomon Algebra As a Free Module For a Generic Arrangement

Nagwan A. Mansoor, Hana M. Ali*

Department of Mathematics, College of Sciences, University of Basrah, Basra, Iraq

*Corresponding author: hana.ali@uobasrah.edu.iq

Doi 10.29072/basjs.20200102

Abstract:

This paper is devoted to study the cohomology of the Orlik-Solomon algebra $A(\mathcal{A})$ as a free module for an ℓ -generic r -arrangement \mathcal{A} and $a = a_i - a_j$, $1 \leq i < j \leq \ell$. In particular, the dimension of $H^k(A(\mathcal{A}), a)$ has been determined for every $1 \leq k \leq r - 1$.

Article inf.

Received

25/12/2019

Accepted

18/2/2020

Published

30/4/2020

Keywords

Hyperplane arrangements,
Hypersolvable arrangements,
Generic arrangements,
Orlik-Solomon algebra



1. Introduction:

The first appearance of the study of hyperplane arrangements was by the researcher S. Robert in (1889, [1]). He defined the arrangement \mathcal{A} as a finite set of lines in the real plane and the number of remaining areas of the plane after deleting those lines was calculated. By a hyperplane H in a finite dimensional vector space V over a field $K = \mathbb{R}$ or \mathbb{C} , we mean an affine subspace of codimension one (that is $\dim H = \dim V - 1$) and by a hyperplane arrangement \mathcal{A} , (or for shorten (an arrangement)) is a finite collection of hyperplanes in V . One of the most essential problems in the topological study in the field of arrangements is the determination of the topological invariants of the complement of an arrangement, $M(\mathcal{A}) = V \setminus \bigcup_{H \in \mathcal{A}} H$, in terms of combinatorics, i.e. those invariants, that can be determined by using the intersection lattice $L(\mathcal{A}) = \{X \subseteq V \mid X = \bigcap_{H \in B} H \text{ and } B \subseteq \mathcal{A}\}$ only, which is partially ordered by: $X \leq Y \Leftrightarrow Y \subseteq X$, that ordered the objects of $L(\mathcal{A})$ opposite of inclusion. As a best general reference here, we refer the reader to [2].

E. Fadell, R. Fox and L. Neuwirth, in (1962, [3] and [4]), studied the cohomological group of the complement of an arrangement in complex space. Perhaps, the first non-trivial result is due to Brieskorn, Orlik and Solomon who calculated the cohomology algebra of the complement in terms of generators and relations (see [5] and [6]). Orlik and Solomon theorem's state that the cohomology algebra of the complement of a complex hyperplane arrangement is isomorphic to an algebra (that named by their names and denoted by O-S algebra) that is combinatorial determined, in line for [6].

This paper is interested with Hattori class of arrangements (1975, [7]) as a subclass of the class of a hypersolvable arrangement which firstly introduced in (1998, [8]) and (2002, [9]) by Jambu and Papadima as a generalization of Stanley class of arrangements (1972, [10]). We emphasize that every hypersolvable arrangement \mathcal{A} has a natural partition on the hyperplanes of \mathcal{A} , we call it a hypersolvable partition and denote it by HP due (2006, [11]),. As well as, a natural ordering was defined on the hyperplanes of \mathcal{A} induced by a fixed HP, we call it a hypersolvable ordering of \mathcal{A} and denoted it by \preceq . The advantage of giving any ℓ – generic r – arrangement \mathcal{A} , a fashion as a hypersolvable arrangement with HP $\Pi = (\Pi_1, \dots, \Pi_\ell)$ and exponent vector $w = (w_1, \dots, w_\ell) = (1, \dots, 1)$, lies in the fact that we can use our knowledge about the structure of the O-S algebra $A(\mathcal{A})$ as a free module with the NBC monomial basis that related to the section of Π .

A subarrangement $C \subseteq \mathcal{A}$ is called a circuit if it is a minimal dependent subarrangement of \mathcal{A} . If H is the smallest hyperplane in C via a total ordering \preceq on the hyperplanes of \mathcal{A} , then $\bar{C} = C \setminus \{H\}$ is called a broken circuit of a circuit C . By NBC base B of \mathcal{A} , we mean a subarrangement of \mathcal{A} which contains no broken circuit and such subarrangements must be independent and denoted by k -NBC base if $|B| = k$. Let \mathcal{A} be an ℓ – generic r – arrangement and for $1 \leq k \leq r$, $NBC_k(\mathcal{A})$ be the set of all k -NBC bases of \mathcal{A} , then;

$$|NBC_k(\mathcal{A})| = \binom{\ell}{k} \text{ and } |NBC_r(\mathcal{A})| = \binom{\ell}{r} - \binom{\ell-1}{r} = \binom{\ell-1}{r-1}, [11, 12].$$



Moreover, for any partition $\Pi = (\Pi_1, \dots, \Pi_\ell)$ on the hyperplanes of an arrangement \mathcal{A} , we recall the definition of the partition \mathcal{K} -module to be $(\Pi)_* = (\Pi_1)_* \otimes \dots \otimes (\Pi_\ell)_*$, where for $1 \leq k \leq \ell$, $(\Pi_k)_*$ is the free \mathcal{K} -module with basis 1 and the elements of Π_k [2]. Similarly, we recall a fashion of the Orlik-Solomon algebra $A(\mathcal{A})$ as a free \mathcal{K} -submodule of a hypersolvable arrangement \mathcal{A} , that was clarified by Al-Taai, Ali and Majeed (2009, [13]).

The study of the Orlik-Solomon algebra $A(\mathcal{A})$ is motivated by creating some topological invariants of the complement $M(\mathcal{A})$ of an arrangement \mathcal{A} . For example, for a real r -arrangement in \mathbb{R}^r , the number of regions of the complement $M(\mathcal{A})$ is, $\sum_{i=1}^r \dim A_i(\mathcal{A})$, see [14]. For a complex r -arrangement \mathcal{A} in \mathbb{C}^r , the complement $M(\mathcal{A})$ is a path-connected manifold which is a minimal CW space for some cases and for $1 \leq k \leq r$, $\dim A_k(\mathcal{A})$ represents the number of the k -cells in each skeleton. Furthermore, for $a \in A_1$, one can define a local coefficient system $\mathcal{L}(a)$ and the connection between $H^*(M(\mathcal{A}), \mathcal{L}(a))$ and the cohomology of the Orlik-Solomon algebra $H^*(A(\mathcal{A}); a)$ has been studied in many papers. In fact, there are many results relating to $\dim H^1(A(\mathcal{A}); a)$. In the case $\text{char } \mathcal{K} = 0$, it has been shown in [15] that $\dim H^1(A(\mathcal{A}); a)$ can be determined by a particular set of elements from $L(\mathcal{A})$. However, little is known about the higher dimension $\dim H^p(A(\mathcal{A}); a)$ for $p > 1$, [15, 16]. This paper contains four sections. The first one is devoted to introduce the preliminaries that we needed in this work. The motivation of each one of the other sections is to calculate the dimension of $H^k(A(\mathcal{A}); a)$ for $1 \leq k \leq r - 1$.

2. PRELIMINARIES

In this section we briefly sketch the notions; O-S algebra, NBC module and partition module. Moreover, we will introduce a definition of the cohomology for O-S algebra. From now on, we will make assumption that $\mathcal{A} = \{H_1, \dots, H_n\}$ is an essential complex r -arrangement with an arbitrary total ordering \preceq on the hyperplanes of it.

2.1 The Orlik-Solomon Algebra: [2]

Let $\{e_{H_1}, \dots, e_{H_n}\}$ be a set that one to one correspondence with \mathcal{A} via the ordering \preceq and let \mathcal{K} be any commutative ring. The Orlik-Solomon algebra (or for shorten O-S algebra) $A_*(\mathcal{A})$ is defined to be the quotient of the exterior \mathcal{K} -algebra, $E_* = \bigwedge_{k \geq 0} (\bigoplus_{H \in \mathcal{A}} K e_H)$, by the homogeneous ideal $I_*(\mathcal{A})$ is generated by the relations, $\sum_{j=1}^k (-1)^{k-1} e_{H_{i_1}} \dots \widehat{e_{H_{i_j}}} \dots e_{H_{i_k}}$, for all $1 \leq i_1 < \dots < i_k \leq n$ such that $\{H_{i_1}, \dots, H_{i_k}\}$ is dependent subarrangement of \mathcal{A} , where the circumflex $\widehat{}$ means $e_{H_{i_j}}$ is deleted.

Assume we have $\partial_*^E: E_* \rightarrow E_*$ to be a K -linear mapping defined as; $\partial_0^E(e_{\emptyset_\ell}) = 0$; $\partial_1^E(e_H) = 1$, for all $H \in \mathcal{A}$ and for $2 \leq k \leq r$, $\partial_k^E(e_C) = \sum_{j=1}^k (-1)^{k-1} e_{H_{i_1}} \dots \widehat{e_{H_{i_j}}} \dots e_{H_{i_k}}$, $C = \{H_{i_1}, \dots, H_{i_k}\}$. Notice that (E_*, ∂_*^E) forms a chain complex, that is called an exterior complex and the chain complex $(A_*(\mathcal{A}), \partial_*^A)$ receives a structure as acyclic chain complex from the exterior complex (E_*, ∂_*^E) , where $\partial_*^A = \psi_* \circ \partial_*^E$ and $\psi_*: E_* \rightarrow A_*(\mathcal{A})$ be the canonical chain map. The acyclic chain complex $(A_*(\mathcal{A}), \partial_*^A)$ is called the O-S complex.



2.2 The NBC module: [2]

For any commutative ring \mathcal{K} , the broken circuit module $NBC_*(\mathcal{A})$ of the exterior \mathcal{K} -algebra E_* defined as; $NBC_0(\mathcal{A}) = \mathcal{K}$ and for $1 \leq k \leq r$; $NBC_k(\mathcal{A})$ is the free \mathcal{K} -module of E_k with (no broken circuit) monomials basis $\{e_C | C \in NBC_k(\mathcal{A})\}$, i.e.;

$$NBC_k(\mathcal{A}) = \bigoplus_{C \in NBC_k(\mathcal{A})} \mathcal{K}e_C \text{ and } NBC_*(\mathcal{A}) = \bigoplus_{k=0}^r NBC_k(\mathcal{A}).$$

Notice that, the broken circuit subcomplex $(NBC_*(\mathcal{A}), \partial_*^{NBC})$ inherits a structure as a cyclic chain complex from the exterior complex (E_*, ∂_*^E) , where $\partial_*^{NBC} = \partial_*^E \circ i_*$ and $i_*: E_* \rightarrow NBC_*(\mathcal{A})$ is the inclusion chain map. Moreover, the restriction of the canonical chain map $\psi_*: E_* \rightarrow A_*(\mathcal{A})$ of the broken circuit module $NBC_*(\mathcal{A})$, is a chain isomorphism, defined as;

$$\text{For } 1 \leq k \leq r, \psi_k(e_C) = e_C + I_k(\mathcal{A}) = a_C, C \in NBC_k(\mathcal{A}).$$

$$\begin{array}{ccccccccccc} 0 \rightarrow & NBC_r(\mathcal{A}) & \xrightarrow{\partial_r^{NBC}} & NBC_{r-1}(\mathcal{A}) & \xrightarrow{\partial_{r-1}^{NBC}} & \dots & \xrightarrow{\partial_2^{NBC}} & NBC_1(\mathcal{A}) & \xrightarrow{\partial_1^{NBC}} & NBC_0(\mathcal{A}) & \xrightarrow{\partial_0^{NBC}} & 0 \\ & \psi_r \downarrow & & \psi_{r-1} \downarrow & & & & \psi_1 \downarrow & & \psi_0 \downarrow & & \\ 0 \rightarrow & A_r(\mathcal{A}) & \xrightarrow{\partial_r^A} & A_{r-1}(\mathcal{A}) & \xrightarrow{\partial_{r-1}^A} & \dots & \xrightarrow{\partial_2^A} & A_1(\mathcal{A}) & \xrightarrow{\partial_1^A} & A_0(\mathcal{A}) & \xrightarrow{\partial_0^A} & 0 \end{array}$$

Thus the O-S algebra has the following structure as a free \mathcal{K} -module:

$$A_*(\mathcal{A}) = \bigoplus_{k=0}^r A_k(\mathcal{A}) = \bigoplus_{k=0}^r (\bigoplus_{C \in NBC_k(\mathcal{A})} \mathcal{K}a_C).$$

We emphasize that, the Poincare polynomials of an arrangement \mathcal{A} and the O-S algebra $A_*(\mathcal{A})$ are equal, i.e. $P(\mathcal{A}, t) = P(A_*(\mathcal{A}), t)$. Therefore, for $1 \leq k \leq r$, the k^{th} Betti number b_k of the Poincare polynomial $P(\mathcal{A}, t)$ will be $b_k = |NBC_k(\mathcal{A})|$.

2.3 Partition module: [2]

Let $\Pi = (\Pi_1, \dots, \Pi_\ell)$ be a partition on an r -arrangement \mathcal{A} and let \mathcal{K} be any commutative ring. A partition \mathcal{K} -module is defined to be $(\Pi)_* = (\Pi_1)_* \otimes \dots \otimes (\Pi_\ell)_*$, where for $1 \leq k \leq \ell$, $(\Pi_k)_*$ is the free \mathcal{K} -module with basis 1 and the elements of Π_k . For each $B = \{H_{i_1}, \dots, H_{i_k}\} \in S_k(\Pi)$ the set of all k -sections of Π , i.e. $H_{i_m} \in \Pi_{i_m}, 1 \leq i_1 < \dots < i_k \leq \ell$ and $1 \leq m \leq k$, define; $q_B = x_1 \otimes \dots \otimes x_\ell \in (\Pi)_*$ as;

$$x_j = \begin{cases} H_j & \text{if } j = i_m \text{ for some } 1 \leq m \leq k \\ 1 & \text{if } j \neq i_m \text{ for all } 1 \leq m \leq k \end{cases}$$

We agree that each of $q_{\emptyset_\ell} = 1 \otimes \dots \otimes 1$ and q_B is homogeneous of degree k . Denote $(\Pi)_k$ the k^{th} -homogeneous part of $(\Pi)_*$. Therefore,

$$(\Pi)_* = \bigoplus_{k=0}^\ell (\Pi)_k = \bigoplus_{k=0}^\ell (\bigoplus_{B \in S_k(\Pi)} \mathcal{K}q_B)$$

and $\{q_B | B \in S_k(\Pi)\}$ forms a basis for the free \mathcal{K} -module $(\Pi)_*$. Furthermore, $\{q_{\{H\}} | H \in \Pi_k\}$ forms a basis for the free \mathcal{K} -module $(\Pi_k)_*, 1 \leq k \leq \ell$. Define a \mathcal{K} -linear mapping $\partial_*^\Pi: (\Pi)_* \rightarrow (\Pi)_*$ as; $\partial_0^\Pi(q_{\{H\}}) = 0, \partial_1^\Pi(q_H) = 1$, for all $H \in \mathcal{A}$ and for $2 \leq k \leq \ell$, $\partial_k^\Pi(q_B) = \sum_{j=1}^k (-1)^{k-1} \widehat{q}_{B_j}$, where $B = \{H_{i_1}, \dots, H_{i_k}\} \in S_k(\Pi), q_B = x_1 \otimes \dots \otimes x_\ell$, and $\widehat{q}_{B_j} = x_1 \otimes \dots \otimes \widehat{H}_{i_j} \otimes \dots \otimes x_\ell$ by means of $\widehat{H}_{i_j} = 1$. ∂_*^π is a differentiation on $(\Pi)_*$ and the chain complex $((\Pi)_*, \partial_*^\pi)$ is called the partition complex. For $1 \leq k \leq \ell$, define the a map

$\tilde{\varphi}_k: \{q_B | B \in S_k(\Pi)\} \rightarrow A_*(\mathcal{A})$, as $\varphi_k(q_B) = a_B = e_B + I_k(\mathcal{A})$, $B \in S_k(\Pi)$. Let $\varphi_k: (\Pi)_k \rightarrow A_k(\mathcal{A})$ be the unique \mathcal{K} -linear map that extend this assignment. Accordingly, there is a unique K -chain mapping $\varphi_*: (\Pi)_* \rightarrow A_*(\mathcal{A})$ between acyclic chain complexes.

2.4 Theorem [13]

Let \mathcal{A} be an $\ell -$ generic $r -$ arrangement. Then;

1. for $1 \leq k \leq r - 1$, be $NBC_k(\mathcal{A}) = S_k(\Pi)$ and $NBC_r(\mathcal{A}) = S_r(\Pi) \setminus (S_r(\Pi) \cap BC_r(\mathcal{A}))$, where $BC_r(\mathcal{A})$ is the set of all $r -$ broken circuits of \mathcal{A} and;
2. The O-S algebra has a fashion as a submodule of the partition module (Π) with, $b_k(A(\mathcal{A})) = |S_k(\Pi)| = \binom{\ell}{k}$ and $b_r(A(\mathcal{A})) = \binom{\ell-1}{r-1}$.

2.5 Cohomology of O-S algebra: [16]

Let $a \in A_1(\mathcal{A})$ with $a = \sum_{i=1}^n \lambda_i a_i$ for $\lambda_i \in \mathcal{K}$. Multiplication by a giving the differential $d_k: A_k(\mathcal{A}) \xrightarrow{a} A_{k+1}(\mathcal{A})$ forms a complex $(A(\mathcal{A}), a)$. The cohomology of this complex is said to be a cohomology of the O-S algebra and it is denoted by $H^*(A(\mathcal{A}), a)$.

2.6 Theorem: [16]

Let \mathcal{A} be a central hyperplane arrangement. Let $a = \sum_{i=1}^n \lambda_i a_i$ for $\lambda_i \in \mathcal{K}$. If $\sum_{i=1}^n \lambda_i a_i \neq 0$, then $H^*(A(\mathcal{A}), a) = 0$.

2: THE STRUCTURE OF $H^1(A(\mathcal{A}), a)$:

From now on we make the assumption that, \mathcal{A} will be an $\ell -$ generic $r -$ arrangement with HP $\Pi = (\Pi_1, \dots, \Pi_\ell)$ and exponent vector $w = (1, \dots, 1)$ and $\ell(\mathcal{A}) = |\mathcal{A}|$.

3.1 Lemma:

If $a = a_1 - a_i$, for $2 \leq i \leq \ell$, then:

1. $d_1(a_1) = d_1(a_i) = a_1 a_i$ and;
2. $\dim(\text{Im } d_1) = \ell - 1$.

Proof:

Firstly, we study the homomorphism, $d_1: A_1(\mathcal{A}) \xrightarrow{a} A_2(\mathcal{A})$. Since \mathcal{A} is $\ell -$ generic $r -$ arrangement, then $E_1 = \langle e_1, e_2, \dots, e_\ell \rangle \cong A_1(\mathcal{A})$.

For 1:

$$d_1(a_1) = a_1 a_1 - a_1 a_i = -a_1 a_i \text{ and;} \\ d_1(a_i) = a_i a_1 - a_i a_i = -a_1 a_i. \text{ So, } d_1(a_1) = d_1(a_i).$$

For 2:

To prove 2, we study $d_1(a_j)$, for $2 \leq j \neq i \leq \ell$. From the definition of $d_1: A_1 \rightarrow A_2$;

$$d_1(a_j) = a_j \cdot a = a_j a_1 - a_j a_i = -a_1 a_j - a_j a_i = \begin{cases} -a_1 a_j - a_j a_i; & 1 < j < i \leq \ell \\ -a_1 a_j + a_i a_j; & 1 < i < j \leq \ell \end{cases} \dots (2.1.1).$$

Notice that, since \mathcal{A} is $\ell -$ generic $r -$ arrangement, then;

$$E_2 = \langle e_{i_1} e_{i_2} \mid 1 \leq i_1 < i_2 \leq \ell \rangle \cong A_2(\mathcal{A}).$$



So, for $1 < i < j \leq \ell$;

$$\partial_2(a_1 a_i a_j) = a_i a_j - a_1 a_j + a_1 a_i \Rightarrow a_i a_j = \partial_2(a_1 a_i a_j) - a_1 a_i + a_1 a_j \dots (2.1.2)$$

Substitute (2.1.1) in (2.1.2), we have;

$$d_1(a_j) = a_1 a_j - \partial_2(a_1 a_i a_j) + a_1 a_i - a_1 a_j = -\partial_2(a_1 a_i a_j) + a_1 a_i.$$

$$\text{Then } \partial_2(a_1 a_i a_j) = d_1(a_j) - a_1 a_i.$$

Since $e_1 e_i e_j$ is an NBC-monomial, hence $d_1(e_j) - e_1 e_i \notin I_2$; So, $d_1(a_j) - a_1 a_i \neq 0_{A_2}$. Then $d_1(a_j) \neq a_1 a_i$ and the number of such cases is $\ell - i$.

Similarly, for $1 < j < i \leq \ell$;

$$\partial_2(a_1 a_j a_i) = a_j a_i - a_1 a_i + a_1 a_j \neq 0_{A_2}. \text{ So,}$$

$$\partial_2(a_1 a_j a_i) = d_1(a_j) + a_1 a_i \neq 0$$

Since $e_1 e_j e_i$ is an NBC-monomial in E_2 hence $\partial_2(e_1 e_j e_i) \notin I_2$, then $d_1(a_j) \neq -a_1 a_i$ and the number of such cases is $i - 1$.

So, the number of monomial in the bases of $Im d_1$ is $\ell - 1$. ■

3.2 Lemma:

If $a = a_i - a_j$ for all $1 \leq i < j \leq \ell$, then;

1. $d_1(a_i) = d_1(a_j) = -a_i a_j$, and;
2. $\dim(Im d_1) = \binom{\ell-1}{1}$.

Proof:

For 1:

$$d_1(a_i) = a_i \cdot a = a_i a_i - a_i a_j = -a_i a_j \text{ and;}$$

$$d_1(a_j) = a_j \cdot a = a_j a_i - a_j a_j = -a_i a_j. \text{ Thus, } d_1(a_i) = d_1(a_j).$$

For 2:

Now, to find the $\dim(Im d_1)$, we looking closely to the following cases;

$$d_1(a_k) = a_k \cdot a = a_k a_i - a_k a_j = \begin{cases} a_k a_i - a_k a_j; & 1 \leq k < i < j \leq \ell \\ -a_i a_k - a_k a_j; & 1 < i < k < j \leq \ell \dots (2.2.1). \\ -a_i a_k + a_j a_k; & 1 < i < j < k \leq \ell \end{cases}$$

Since $E_2 = \langle e_{i_1} e_{i_2} \mid 1 \leq i_1 < i_2 \leq \ell \rangle \cong A_2(\mathcal{A})$, hence $d_1(a_k)$ is written as a linear combination of NBC monomials. By studying all the cases above, we pointing that there are no $1 \leq k < l \leq \ell$ such that $d_1(a_k) = d_1(a_l)$ and by simple calculation of the number of the generators of $Im(d_1)$ we have $\dim(Im d_1) = \binom{\ell-1}{1}$. ■

3.3 Proposition:

If $a = a_1 - a_j$, $2 \leq j \leq \ell$, then:

1. $d_2(a_1 a_j) = 0_{A_3}$;
2. $\dim(\ker d_2) = \dim(Im d_1)$;
3. $H^1(A(\mathcal{A}); a) = 0$;



4. If $r > 3$, then $\dim(\text{Im } d_2) = \binom{\ell}{2} - \binom{\ell-2}{1}$ and;
5. If $r = 3$, then $\dim(\text{Im } d_2) = \binom{\ell-2}{1}$.

Proof:

Firstly, we will study the homomorphism $d_2: A_2 \rightarrow A_3$ that is defined as;

$$d_2(a_{k_1} a_{k_2}) = a_{k_1} a_{k_2} a_1 - a_{k_1} a_{k_2} a_j, \text{ where } 2 \leq j \leq \ell \text{ and } 1 \leq k_1 < k_2 \leq \ell$$

We have many cases to study as follows;

Case (1):- For $k_1 = 1$ and $k_2 = j$,

$$d_2(a_1 a_j) = a_1 a_j a_1 - a_1 a_j a_j = 0_{A_3}. \text{ Thus, } a_1 a_j \in \ker d_2.$$

Case (2):- If $k_1 = 1$ and $1 < k_2 < j < \ell$, thus;

$$d_2(a_1 a_{k_2}) = a_1 a_{k_2} a_1 - a_1 a_{k_2} a_j = -a_1 a_{k_2} a_j.$$

$$\text{As well as; } d_2(a_{k_2} a_j) = a_{k_2} a_j a_1 - a_{k_2} a_j a_j = a_1 a_{k_2} a_j,$$

$$\text{i.e. } d_2(a_1 a_{k_2}) = -d_2(a_{k_2} a_j) \neq 0_{A_3}.$$

Case (3):- If $k_1 = 1$ and $1 < j < k_2 < \ell$, then;

$$d_2(a_1 a_{k_2}) = a_1 a_j a_{k_2}. \text{ Consequently, } d_2(a_j a_{k_2}) = a_1 a_j a_{k_2}.$$

$$\text{So, } d_2(a_1 a_{k_2}) = d_2(a_j a_{k_2}) \neq 0_{A_3}.$$

Case(4):- If $1 < k_1 < k_2 < j \leq \ell$, then $d_2(a_{k_1} a_{k_2}) = a_1 a_{k_1} a_{k_2} - a_{k_1} a_{k_2} a_j \dots(2.3.1)$.

Since \mathcal{A} is ℓ –generic r –arrangement, here we have two possible cases in this case, are listed below:

- If $r > 3$, we have $d_2(a_{k_1} a_{k_2}) = a_1 a_{k_1} a_{k_2} - a_{k_1} a_{k_2} a_j \neq 0_{A_3}$ is written as a linear combination of NBC- monomials, or;
- If $r = 3$, we have $\partial_4(e_1 e_{k_1} e_{k_2} e_j) \in I_3$. Thus; $a_{k_1} a_{k_2} a_j - a_1 a_{k_2} a_j + a_1 a_{k_1} a_j - a_1 a_{k_1} a_{k_2} = 0_{A_3}$. So, $a_{k_1} a_{k_2} a_j = a_1 a_{k_2} a_j - a_1 a_{k_1} a_j + a_1 a_{k_1} a_{k_2} \dots (2.3.2)$. By substituting (2.3.2) in (2.3.1), we have $d_2(a_{k_1} a_{k_2}) = a_1 a_{k_1} a_{k_2} - a_1 a_{k_2} a_j + a_1 a_{k_1} a_j - a_1 a_{k_1} a_{k_2} = a_1 a_{k_1} a_j - a_1 a_{k_2} a_j \neq 0_{A_3}$ is written as a linear combination of NBC- monomials.

Case (5):- If $1 < k_1 < j < k_2 \leq \ell$, then $d_2(a_{k_1} a_{k_2}) = a_1 a_{k_1} a_{k_2} + a_{k_1} a_j a_{k_2} \dots (2.3.3)$, in this case there are two possible cases as follows;

- If $r > 3$, then $e_{k_1} e_{k_2} e_j$ is an NBC-monomial of E_3 , since $\{H_{k_1}, H_{k_2}, H_j\}$ is an NBC base of \mathcal{A} . Thus; $d_2(a_{k_1} a_{k_2}) = a_1 a_{k_1} a_{k_2} + a_{k_1} a_j a_{k_2} \neq 0_{A_3}$ is written as a linear combination of NBC- monomials, or;
- If $r = 3$, we have $e_{k_1} e_{k_2} e_j$ is a broken circuit monomial of E_3 , since $\{H_{k_1}, H_{k_2}, H_j\}$ is a broken circuit of \mathcal{A} . We know that, $\partial_4(e_1 e_{k_1} e_j e_{k_2}) \in I_3$. Thus;

$$a_{k_1} a_j a_{k_2} - a_1 a_j a_{k_2} + a_1 a_{k_1} a_{k_2} - a_1 a_{k_1} a_j = 0_{A_3} \dots (2.3.4).$$

By substituting (2.3.4) in (2.3.3), we have;

$$d_2(a_{k_1} a_{k_2}) = a_1 a_{k_1} a_{k_2} + a_1 a_j a_{k_2} - a_1 a_{k_1} a_{k_2} + a_1 a_{k_1} a_j = a_1 a_{k_1} a_j + a_1 a_j a_{k_2} \neq 0_{A_3};$$

is written as a linear combination of NBC- monomials.

Case (6):- If $1 < j < k_1 < k_2 < j \leq \ell$, then;



$$d_2(a_{k_1}a_{k_2}) = a_1a_{k_1}a_{k_2} - a_ja_{k_1}a_{k_2} \dots (2.3.5),$$

We have two possible cases can be obtained from this case;

- If $r > 3$, we have $e_j e_{k_1} e_{k_2}$ is an *NBC* monomial of E_3 , since $\{H_j, H_{k_1}, H_{k_2}\}$ is an *NBC* base of \mathcal{A} . Thus, $d_2(a_{k_1}a_{k_2}) = a_1a_{k_1}a_{k_2} - a_ja_{k_1}a_{k_2} \neq 0_{A_3}$ is written as a linear combination of *NBC*- monomials, or;
- If $r = 3$, then $e_j e_{k_1} e_{k_2}$ is a broken circuit monomial of E_3 , since $\{H_j, H_{k_1}, H_{k_2}\}$ is a broken circuit of \mathcal{A} . Thus; $\partial_4(e_1 e_j e_{k_1} e_{k_2}) \in I_3$ and;

$$a_j a_{k_1} a_{k_2} - a_1 a_{k_1} a_{k_2} + a_1 a_j a_{k_2} - a_1 a_j a_{k_1} = 0_{A_3} \dots (2.3.6)$$

By substituting (2.3.6) in (2.3.5), we have, $d_2(a_{k_1}a_{k_2}) = -a_1 a_j a_{k_2} + a_1 a_j a_{k_1}$ is written as a linear combination of *NBC*- monomials.

Now, we clarify our assertion:

For 1:- It is clear from case (1).

For 2 and 3:- From our study of the 6th cases that we discussed above, we have just one 2 – *NBC*-monomial $a_1 a_j$ that satisfied; $d_2(a_1 a_j) = 0_{A_3}$. But $a_1 a_j \in \text{Im } d_1$ and since $\text{Im } d_1 \subseteq \ker d_2$, hence $\text{Im } d_1 = \ker d_2$. Which implies that $H^1(A(\mathcal{A}); a) = 0$.

For 4:- The number of the images of the *NBC* monomials that are need to remove it from the $\binom{\ell}{2}$ – *NBC* monomial came from case (1) and the repetition that came from cases (2) and (3) and the number of such cases is equal to $\ell - 2$. Thus;

$$\dim(\text{Im } d_2) = \binom{\ell}{2} - \ell + 2 = \binom{\ell}{2} - \binom{\ell-2}{1}.$$

For 5:- If $r = 3$, then $\dim(\text{Im } d_2) = \ell - 2 = \binom{\ell-2}{1}$ is the number of the 2 – *NBC* monomial that begin with a_1 and contain no a_j . ■

3.4 Proposition:

If $a = a_i - a_j, 1 < i < j < \ell$, then:-

1. $d_2(a_i a_j) = 0_{A_3}$;
2. $\ker d_2 = \text{Im } d_1$;
3. $H^1(A(\mathcal{A}); a) = 0$;
4. If $r > 3$, $\dim(\text{Im } d_2) = \binom{\ell}{2} - \binom{\ell-2}{1}$ and;
5. If $r = 3$, $\dim(\text{Im } d_2) = \binom{\ell-2}{1}$.

Proof:

We will study the homomorphism $d_2: A_2 \rightarrow A_3$ to serve our aim, as; $d_2(a_{k_1}a_{k_2}) = a_{k_1}a_{k_2}a_i - a_{k_1}a_{k_2}a_j$; where $1 \leq k_1 < k_2 \leq \ell$. We have the following possible cases for choosing k_1 and k_2 ;

Case (1):- If $k_1 = i$ and $k_2 = j$, then $d_2(a_i a_j) = a_i a_j a_i - a_i a_j a_j = 0_{A_3}$. Thus, $a_i a_j \in \ker d_2$. In fact $a_i a_j \in \text{Im } d_1$.

Case (2):- If $k_1 = i$ and $k_2 \neq j$ and $1 \leq k_2 \leq \ell$ then;



$$d_2(a_i a_{k_2}) = a_i a_{k_2} a_i - a_i a_{k_2} a_j = \begin{cases} a_{k_2} a_i a_j; & k_2 < i < j \\ -a_i a_{k_2} a_j; & i < k_2 < j \\ a_i a_j a_{k_2}; & i < j < k_2 \end{cases} = \begin{cases} -d_2(a_{k_2} a_j); & k_2 < i < j \\ -d_2(a_{k_2} a_j); & i < k_2 < j \\ -d_2(a_j a_{k_2}); & i < j < k_2 \end{cases}.$$

We have two possible cases can be obtained from this case;

- If $r > 3$, then $d_2(a_i a_{k_2})$ is written as a linear combination of 3-NBC monomials.
- If $r = 3$, as we know, either $\partial_4(e_1 e_{k_2} e_i e_j) \in I_3$, $\partial_4(e_1 e_i e_{k_2} e_j) \in I_3$ or $\partial_4(e_1 e_i e_j e_{k_2}) \in I_3$ depends of $k_2 < i < j$ or $i < k_2 < j$ or $i < j < k_2$ respectively. So;

$$\begin{aligned} & \text{either, } a_{k_2} a_i a_j - a_1 a_i a_j + a_1 a_{k_2} a_j - a_1 a_{k_2} a_i = 0_{A_3} \text{ or;} \\ & a_i a_{k_2} a_j - a_1 a_{k_2} a_j + a_1 a_i a_j - a_1 a_i a_{k_2} = 0_{A_3} \text{ or;} \\ & a_i a_j a_{k_2} - a_1 a_j a_{k_2} + a_1 a_i a_{k_2} - a_1 a_i a_j = 0_{A_3}. \text{ Thus;} \\ d_2(a_i a_{k_2}) &= \begin{cases} a_1 a_i a_j - a_1 a_{k_2} a_j + a_1 a_{k_2} a_i; & k_2 < i < j \\ -a_1 a_{k_2} a_j + a_1 a_i a_j - a_1 a_i a_{k_2}; & i < k_2 < j \\ a_1 a_j a_{k_2} - a_1 a_i a_{k_2} - a_1 a_i a_j; & i < j < k_2 \end{cases} \\ &= \begin{cases} -d_2(a_1 a_{k_2}) - d_2(a_1 a_i); & k_2 < i < j \\ d_2(a_1 a_{k_2}) + d_2(a_1 a_i); & i < k_2 < j \\ d_2(a_1 a_{k_2}) + d_2(a_1 a_i); & i < j < k_2 \end{cases} \end{aligned}$$

is written as a linear combination of 3-NBC monomials.

Case (4):- If $k_1 \neq i$ or j , $k_2 \neq i$ or j and $1 \leq k_1 < k_2 \leq \ell$, we have;

$$d_2(a_{k_1} a_{k_2}) = a_{k_1} a_{k_2} a_i - a_{k_1} a_{k_2} a_j = \begin{cases} a_{k_1} a_{k_2} a_i - a_{k_1} a_{k_2} a_j; & k_1 < k_2 < i < j \\ -a_{k_1} a_i a_{k_2} - a_{k_1} a_{k_2} a_j; & k_1 < i < k_2 < j \\ a_i a_{k_1} a_{k_2} - a_{k_1} a_{k_2} a_j; & i < k_1 < k_2 < j \dots \\ a_i a_{k_1} a_{k_2} + a_{k_1} a_j a_{k_2}; & i < k_1 < j < k_2 \\ a_i a_{k_1} a_{k_2} - a_j a_{k_1} a_{k_2}; & i < j < k_1 < k_2 \end{cases}$$

(2.4.1)

If $r > 3$, then any 3 –monomial $a_{i_1} a_{i_2} a_{i_3}$ is an NBC monomial and $d_2(a_{k_1} a_{k_2})$ is written as a linear combination of 3-NBC monomials.

- If $r = 3$, we have $\partial_4(a_1 a_{k_1} a_{k_2} a_j) = 0_{A_3}$, since $\{H_1, H_{k_1}, H_{k_2}, H_j\}$ is dependent. So, $a_{k_1} a_{k_2} a_j - a_1 a_{k_2} a_j + a_1 a_{k_1} a_j - a_1 a_{k_1} a_{k_2} = 0_{A_3} \dots$ (2.4.2). So we have the following cases:-

Case (4.a):- If $k_1 < k_2 < i$, then $\partial_4(a_1 a_{k_1} a_{k_2} a_i) = 0_{A_3}$, since $\{H_1, H_{k_1}, H_{k_2}, H_i\}$ is dependent. So $a_{k_1} a_{k_2} a_i - a_1 a_{k_2} a_i + a_1 a_{k_1} a_i - a_1 a_{k_1} a_{k_2} = 0_{A_3} \dots$ (2.4.3). From the equations (2.4.1), (2.4.2) and (2.4.3) above we have;

$$d_2(a_{k_1} a_{k_2}) = a_1 a_{k_2} a_i - a_1 a_{k_1} a_i + a_1 a_{k_1} a_{k_2} - a_1 a_{k_2} a_j + a_1 a_{k_1} a_j - a_1 a_{k_1} a_{k_2} = -d_2(a_1 a_{k_1}) + d_2(a_1 a_{k_2}).$$

Case (4.b):- If $k_1 < i < k_2$, then;

$$\partial_4(a_1 a_{k_1} a_i a_{k_2}) = a_{k_1} a_i a_{k_2} - a_1 a_i a_{k_2} + a_1 a_{k_1} a_{k_2} - a_1 a_{k_1} a_i = 0_{A_3} \dots$$
 (2.4.4).



Since $\{H_1, H_{k_1}, H_i, H_{k_2}\}$ is dependent. From the equations (2.4.1), (2.4.2) and (2.4.4) we have;

$$d_2(a_{k_1} a_{k_2}) = a_{k_1} a_i a_{k_2} - a_{k_1} a_{k_2} a_j = -a_1 a_i a_{k_2} + a_1 a_{k_1} a_{k_2} - a_1 a_{k_1} a_i - a_{k_1} a_{k_2} a_j + a_1 a_{k_1} a_j - a_1 a_{k_1} a_{k_2} = -d_2(a_1 a_{k_1}) + d_2(a_1 a_{k_2}).$$

Case (4.c):- If $i < k_1 < k_2$, then,

$$\partial_4(a_1 a_i a_{k_1} a_{k_2}) = a_i a_{k_1} a_{k_2} - a_1 a_{k_1} a_{k_2} + a_1 a_i a_{k_2} - a_1 a_i a_{k_1} = 0_{A_3} \dots (2.4.5).$$

Since $\{H_1, H_i, H_{k_1}, H_{k_2}\}$ is dependent subarrangement. From the equations (2.4.1), (2.4.2) and (2.4.5) we have;

$$\begin{aligned} d_2(a_{k_1} a_{k_2}) &= a_i a_{k_1} a_{k_2} - a_{k_1} a_{k_2} a_j \\ &= a_1 a_{k_1} a_{k_2} - a_1 a_i a_{k_2} + a_1 a_i a_{k_1} - a_1 a_{k_2} a_j + a_1 a_{k_1} a_j - a_1 a_{k_1} a_{k_2} \\ &= a_1 a_i a_{k_1} + a_1 a_{k_1} a_j - a_1 a_i a_{k_2} - a_1 a_{k_2} a_j = d_2(a_1 a_{k_1}) + d_2(a_1 a_{k_2}). \end{aligned}$$

- If $r = 3$ and $k_1 = 1$, then $d_2(a_1 a_{k_2}) = \begin{cases} a_1 a_{k_2} a_i - a_1 a_{k_2} a_j & \text{if } 1 < i < k_2 < j \\ a_1 a_{k_2} a_i + a_1 a_j a_{k_2} & \text{if } 1 < i < j < k_2 \end{cases}$

written as a linear combination of *NBC*- monomials.

Case (5):-

- If $r > 3$, we have

$$d_2(a_{k_1} a_{k_2}) = a_{k_1} a_{k_2} a_i - a_{k_1} a_{k_2} a_j = \begin{cases} a_i a_{k_1} a_{k_2} + a_{k_1} a_j a_{k_2} & i < k_1 < j < k_2 \\ a_i a_{k_1} a_{k_2} - a_j a_{k_1} a_{k_2} & i < j < k_1 < k_2 \end{cases} \dots (2.4.6)$$

- If $r = 3$, we need to write $d_2(a_{k_1} a_{k_2})$ as a linear combination of *NBC* monomials.

Since $\{H_1, H_i, H_{k_1}, H_{k_2}\}$ is dependent, then;

$$\partial_4(a_1 a_i a_{k_1} a_{k_2}) = a_i a_{k_1} a_{k_2} - a_1 a_{k_1} a_{k_2} + a_1 a_i a_{k_2} - a_1 a_i a_{k_1} = 0_{A_3} \dots (2.4.7),$$

we have two cases as;

Case (5.1):- If $k_1 < j < k_2$, then $\partial_4(a_1 a_i a_{k_1} a_{k_2}) = 0_{A_3}$ since $\{H_1, H_{k_1}, H_j, H_{k_2}\}$ is dependent. Thus, $a_{k_1} a_j a_{k_2} - a_1 a_j a_{k_2} + a_1 a_{k_1} a_{k_2} - a_1 a_{k_1} a_j = 0_{A_3} \dots (2.4.8).$

From the above equations (2.4.6), (2.4.7) and (2.4.8) we have;

$$\begin{aligned} d_2(a_{k_1} a_{k_2}) &= a_1 a_{k_1} a_{k_2} - a_1 a_i a_{k_2} + a_1 a_i a_{k_1} + a_1 a_j a_{k_2} - a_1 a_{k_1} a_{k_2} + a_1 a_{k_1} a_j \\ &= a_1 a_i a_{k_1} + a_1 a_{k_1} a_j - a_1 a_i a_{k_2} + a_1 a_j a_{k_2} = -d_2(a_1 a_{k_1}) + d_2(a_1 a_{k_2}). \end{aligned}$$

Case (5.2):- If $j < k_1 < k_2$, then $\partial_4(a_1 a_j a_{k_1} a_{k_2}) = 0_{A_3} \dots (2.4.9)$, since $\{H_1, H_j, H_{k_1}, H_{k_2}\}$ is dependent subarrangement. Thus, we have;

$$\begin{aligned} d_2(a_{k_1} a_{k_2}) &= a_1 a_{k_1} a_{k_2} - a_1 a_i a_{k_2} + a_1 a_i a_{k_1} - a_1 a_{k_1} a_{k_2} + a_1 a_j a_{k_2} - a_1 a_j a_{k_1} \\ &= a_1 a_i a_{k_1} - a_1 a_j a_{k_1} - a_1 a_i a_{k_2} + a_1 a_j a_{k_2} = -a_1 a_{k_1} a_i + a_1 a_{k_1} a_j + a_1 a_{k_2} a_i - a_1 a_{k_2} a_j \\ &= -d_2(a_1 a_{k_1}) + d_2(a_1 a_{k_2}). \end{aligned}$$

Now, we verify our claim as follows:-

For 1:- See case 1.

For 2:- From case(1), we have just one $\ell - NBC$ monomial satisfied $d_2(a_i a_j) = 0_{A_3}$, which is a coboundary. Since $a_i a_j \in Im d_1 \subseteq ker d_2$, hence $ker d_2 = Im d_1$.

For 3:- It is clear that, $H^1(A(\mathcal{A}); a) = 0$.



For 4:- If $r > 3$, the repetition that we discussed them in the cases 2 and 3 and from case 1 is $\ell - 2$; $\dim(\text{Im } d_2) = \binom{\ell}{2} - \ell + 2$.

For 5:- If $r = 3$, from our discussion of all cases. If $r = 3$, it is clear that the collection $B = \{d_2(a_1 a_{k_2}) : 2 \leq k_2 \leq \ell \text{ and } k_2 \neq i \text{ or } j\} \cup \{d_2(a_1 a_{k_1})\}$ play as a basis for $\text{Im } d_2$, so; $\dim d_2 = |B| = (\ell - 3) + 1 = \ell - 2 = \binom{\ell-2}{1}$. ■

4.THE STRUCTURE OF $H^2(A(\mathcal{A}), a)$ AND $H^3(A(\mathcal{A}), a)$:

4.1 Corollary:

If $r = 3$ and $a = a_1 - a_j$ for $2 \leq j \leq \ell$; then $\dim H^2(A(\mathcal{A}), a) = \binom{\ell-1}{2} - \binom{\ell-2}{1}$.

Proof:

Since $d_3: A_3 \rightarrow 0$ is the zero homomorphism. Then $\ker d_3 = A_3$ and $\dim \ker d_3 = \binom{\ell-1}{2}$. Since $\text{Im } d_2 \subseteq \ker d_3$ and from proposition (2.3), we have;

$$\begin{aligned} \dim H^2(A(\mathcal{A}), a) &= \dim \ker d_3 - \dim \text{Im } d_2 \\ &= \dim A_3 - \dim \text{Im } d_2 = \binom{\ell-1}{2} - \ell + 2 = \binom{\ell-1}{2} - \binom{\ell-2}{1}. \blacksquare \end{aligned}$$

4.2 Corollary:

If $r = 3$ and $a = a_i - a_j$ for $1 < i < j \leq \ell$, we have $\dim H^2(A(\mathcal{A}), a) = \binom{\ell-1}{2} - \binom{\ell-2}{1}$.

Proof:

Since $d_3: A_3 \rightarrow 0$ is the zero homomorphism, then $\ker d_3 = A_3$ and since $\text{Im } d_2 \subseteq \ker d_3$, then;

$$\begin{aligned} \dim H^2(A(\mathcal{A}), a) &= \dim(\ker d_3 / \text{Im } d_2) \\ &= \dim A_3 - \dim \text{Im } d_2 = \binom{\ell-1}{2} - \ell + 2 = \binom{\ell-1}{2} - \binom{\ell-2}{1} \blacksquare \end{aligned}$$

4.3 Proposition:

If $r > 3$ and $a = a_1 - a_j$, $2 \leq j \leq \ell$. Then:-

1. $d_3(a_1 a_m a_j) = d_3(a_1 a_j a_m) = 0_{A_4}$, if either $1 < m < j \leq \ell$ or $1 < j < m \leq \ell$.
2. $\ker d_3 = \text{Im } d_2$.
3. $H^2(A(\mathcal{A}), a) = 0$.
4. If $r > 4$, $\dim(\text{Im } d_3) = \binom{\ell}{3} - \binom{\ell-2}{2}$.
5. If $r = 4$, $\dim(\text{Im } d_3) = \binom{\ell}{3} - \binom{\ell-2}{2}$.

Proof:

Firstly, we will study the homomorphism $d_3: A_3 \rightarrow A_4$. As we know the NBC-monomial basis for A_3 is; $B_3 = \{a_{m_1} a_{m_2} a_{m_3} : \{H_{m_1}, H_{m_2}, H_{m_3}\} \subseteq \mathcal{A} \text{ and } 1 \leq m_1 < m_2 < m_3 \leq \ell\}$ and that will enable us to exercise d_3 . So, $d_3(a_{m_1} a_{m_2} a_{m_3}) = a_{m_1} a_{m_2} a_{m_3} a_i - a_{m_1} a_{m_2} a_{m_3} a_j$. To serve our aim we will discuss all the possible cases.

Case (1):- If $m_1 = 1$ and m_2 or $m_3 = j$. Then as we mentioned it previously, $d_3(a_1 a_j a_{m_3}) = d_3(a_1 a_{m_2} a_j) = 0_{A_4}$.

Case (2):- If $m_1 = 1$ and $m_2, m_3 \neq j$ then;



$$d_3(a_1 a_{m_2} a_{m_3}) = a_1 a_{m_2} a_{m_3} a_i = \begin{cases} a_1 a_{m_2} a_{m_3} a_j & m_2 < m_3 < j \leq \ell \\ -a_1 a_{m_2} a_j a_{m_3} & m_2 < j < m_3 \leq \ell \\ a_1 a_j a_{m_2} a_{m_3} & j < m_2 < m_3 \leq \ell \end{cases}$$

$$= \begin{cases} -d_3(a_{m_2} a_{m_3} a_j) & m_2 < m_3 < j \leq \ell \\ d_3(a_{m_2} a_j a_{m_3}) & m_2 < j < m_3 \leq \ell \\ -d_3(a_j a_{m_2} a_{m_3}) & j < m_2 < m_3 \leq \ell \end{cases} \neq 0_{A_4}.$$

Notice that $d_3(a_1 a_{m_2} a_{m_3})$ is an 4 – NBC monomial for any $r \geq 4$. The number of such repetitions is equal to the number of the 3 – NBC monomials that begin with a_1 and the number of such monomials equal to the number of 2 – NBC monomials that not begin with 1 and not equal to j. Thus we have $\binom{\ell-2}{2}$ repetition in this case.

Case (3):- If $1 \leq m_1 < m_2 < m_3 \leq \ell$ and $m_n \neq i$ or j for $n = 1, 2, 3$. Then we have the following cases:-

Case (3.a):- If $1 < m_1 < m_2 < m_3 < j \leq \ell$, then;

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = a_{m_1} a_{m_2} a_{m_3} a_1 - a_{m_1} a_{m_2} a_{m_3} a_j$$

$$= -a_1 a_{m_1} a_{m_2} a_{m_3} - a_{m_1} a_{m_2} a_{m_3} a_j \dots (3.3.1)$$

- If $r > 4$, then $d_3(a_{m_1} a_{m_2} a_{m_3}) \neq 0_{A_4}$ can be written as a linear combination of 4 – NBC monomials.
- If $r = 4$, then we have $\partial_5(e_1 e_{m_1} e_{m_2} e_{m_3} e_j) \in I_4$; $a_{m_1} a_{m_2} a_{m_3} a_j - a_1 a_{m_2} a_{m_3} a_j + a_1 a_{m_1} a_{m_3} a_j - a_1 a_{m_1} a_{m_2} a_j + a_1 a_{m_1} a_{m_2} a_{m_3} \neq 0_{A_4} \dots (3.3.2)$

From the equations (3.3.1) and (3.3.2) above, we have;

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = -a_1 a_{m_1} a_{m_2} a_{m_3} - a_1 a_{m_2} a_{m_3} a_j + a_1 a_{m_1} a_{m_3} a_j$$

$$- a_1 a_{m_1} a_{m_2} a_j + a_1 a_{m_1} a_{m_2} a_{m_3} = -a_1 a_{m_2} a_{m_3} a_j + a_1 a_{m_1} a_{m_3} a_j - a_1 a_{m_1} a_{m_2} a_j$$

$$= -d_3(a_1 a_{m_2} a_{m_3}) + d_3(a_1 a_{m_1} a_{m_3}) - d_3(a_1 a_{m_1} a_{m_2}) \neq 0_{A_4}.$$

Case (3.b):- We have three cases:

Case (3.b.1):- If $1 < m_1 < m_2 < j < m_3 \leq \ell$; then $d_3(a_{m_1} a_{m_2} a_{m_3}) = a_{m_1} a_{m_2} a_{m_3} a_1 - a_{m_1} a_{m_2} a_{m_3} a_j = -a_1 a_{m_1} a_{m_2} a_{m_3} + a_{m_1} a_{m_2} a_j a_{m_3} \dots (3.3.3)$.

- If $r > 3$, then $d_3(a_{m_1} a_{m_2} a_{m_3}) \neq 0_{A_4}$ can be written as a linear combination of 4 – NBC – monomials.
- If $r = 3$, then we have $\partial_5(e_1 e_{m_1} e_{m_2} e_j e_{m_3}) \in I_4$. So;

$$a_{m_1} a_{m_2} a_j a_{m_3} - a_1 a_{m_2} a_j a_{m_3} + a_1 a_{m_1} a_j a_{m_3} - a_1 a_{m_1} a_{m_2} a_{m_3} + a_1 a_{m_1} a_{m_2} a_j = 0_{A_4} \dots$$

(3.3.4). From (3.3.3) and (3.3.4) we have;

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = -a_1 a_{m_1} a_{m_2} a_{m_3} + a_1 a_{m_2} a_j a_{m_3} - a_1 a_{m_1} a_j a_{m_3} + a_1 a_{m_1} a_{m_2} a_{m_3} - a_1 a_{m_1} a_{m_2} a_j$$

$$= a_1 a_{m_2} a_j a_{m_3} - a_1 a_{m_1} a_j a_{m_3} - a_1 a_{m_1} a_{m_2} a_j = -d_3(a_1 a_{m_2} a_{m_3}) + d_3(a_1 a_{m_1} a_{m_3}) - d_3(a_1 a_{m_1} a_{m_2}).$$

Case (3.b.2):- If $1 < m_1 < j < m_2 < m_3 \leq \ell$, then;

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = a_{m_1} a_{m_2} a_{m_3} a_1 - a_{m_1} a_{m_2} a_{m_3} a_j = -a_1 a_{m_1} a_{m_2} a_{m_3} - a_{m_1} a_j a_{m_2} a_{m_3} \dots$$

(3.3.5).



- If $r > 3$, then $d_3(a_{m_1}a_{m_2}a_{m_3}) \neq 0_{A_4}$ written as a linear combination of the 4 – NBC- monomials.
- If $r = 3$, then we have $\partial_5(e_1e_{m_1}e_je_{m_2}e_{m_3}) \in I_4$. So;

$$a_{m_1}a_ja_{m_2}a_{m_3} - a_1a_ja_{m_2}a_{m_3} + a_1a_{m_1}a_{m_2}a_{m_3} - a_1a_{m_1}a_ja_{m_3} + a_1a_{m_1}a_ja_{m_2} = 0_{A_4} \dots$$

(3.3.6). From (3.3.5) and (3.3.6), we have;

$$d_3(a_{m_1}a_{m_2}a_{m_3}) = -a_1a_{m_1}a_{m_2}a_{m_3} - a_1a_ja_{m_2}a_{m_3} + a_1a_{m_1}a_{m_2}a_{m_3} - a_1a_{m_1}a_ja_{m_3} + a_1a_{m_1}a_ja_{m_2} = -a_1a_ja_{m_2}a_{m_3} - a_1a_{m_1}a_ja_{m_3} + a_1a_{m_1}a_ja_{m_2} = -d_3(a_1a_{m_2}a_{m_3}) + d_3(a_1a_{m_1}a_{m_3}) - d_3(a_1a_{m_1}a_{m_2}) \neq 0_{A_4}.$$

Case (3.b.3):- If $1 < j < m_1 < m_2 < m_3 \leq \ell$, then $d_3(a_{m_1}a_{m_2}a_{m_3}) = a_{m_1}a_{m_2}a_{m_3}a_1 - a_{m_1}a_{m_2}a_{m_3}a_j = -a_1a_{m_1}a_{m_2}a_{m_3} + a_ja_{m_1}a_{m_2}a_{m_3} \dots$ (3.3.7).

- If $r > 3$, $d_3(a_{m_1}a_{m_2}a_{m_3}) \neq 0_{A_4}$ can be written as a linear combination of the 4 – NBC monomials.
- If $r = 3$, then we have $\partial_5(e_1e_je_{m_1}e_{m_2}e_{m_3}) \in I_4$. So;

$$a_ja_{m_1}a_{m_2}a_{m_3} - a_1a_{m_1}a_{m_2}a_{m_3} + a_1a_ja_{m_2}a_{m_3} - a_1a_ja_{m_1}a_{m_3} + a_1a_ja_{m_1}a_{m_2} = 0_{A_4} \dots$$

(3.3.8). From (3.3.7) and (3.3.8), we have;

$$d_3(a_{m_1}a_{m_2}a_{m_3}) = -a_1a_{m_1}a_{m_2}a_{m_3} + a_1a_{m_1}a_{m_2}a_{m_3} - a_1a_ja_{m_2}a_{m_3} + a_1a_ja_{m_1}a_{m_3} + a_1a_ja_{m_1}a_{m_2} = -a_1a_ja_{m_2}a_{m_3} + a_1a_ja_{m_1}a_{m_3} - a_1a_ja_{m_1}a_{m_2} = -d_3(a_1a_{m_2}a_{m_3}) + d_3(a_1a_{m_1}a_{m_3}) - d_3(a_1a_{m_1}a_{m_2}) \neq 0_{A_4}.$$

To prove our claim;

For (1): see case (1).

For (2): From case (1) above and lemma (3), we have $a_1a_ma_ja, a_1a_ja_n \in \ker d_3$ and $a_1a_ma_ja, a_1a_ja_n \in \text{Im } d_2$ for all $11 < m < j < n \leq \ell$. Since $\text{Im } d_2 \subseteq \ker d_3$, so $\text{Im } d_2 = \ker d_3$.

For (3): Our claim is done from case (2); i.e. $H^2(A(\mathcal{A}), a) = 0$.

For (4): If $r > 4$, then from all cases that studied in our proof; we have $\dim \text{Im } d_3 = \binom{\ell}{3} - \left(\binom{\ell}{2} + 2\ell - 3\right) = \binom{\ell}{3} \binom{\ell-2}{2}$.

For (5): If $r = 4$, then from all the cases we studied above we have $\dim \text{Im } d_3 = \binom{\ell}{2} - (\ell - 1) - (\ell - 2) = \binom{\ell}{2} - 2\ell + 3 = \binom{\ell-2}{2}$, which is depend on 4 – NBC bases that begins by a_1 that equal to the number 2 – NBC bases which is not contains the number of 3 – NBC- monomial that it contained in $\ker d_3$, that is $d_3(a_1a_ja_m) = d_3(a_1a_ma_ja) = 0_{A_4}$. ■

4.4 Corollary :

If $r = 4$ and $a = a_1 - a_j, 2 \leq j \leq \ell$. Then, $\dim H^3(A(\mathcal{A}), a) = \binom{\ell-1}{3} - \binom{\ell-2}{2}$.

Proof:

Since $d_4: A_4 \rightarrow 0$ is the zero homomorphism, then $\ker d_4 = A_4$ and $\dim \ker d_4 = \binom{\ell-1}{3}$.

By applying Proposition (3.1), we have;



$$\dim H^3(A(\mathcal{A}), a) = \dim(\ker d_4 / \text{Im } d_3) = \dim(\ker d_4) - \dim(\text{Im } d_3) = \binom{\ell-1}{3} - \binom{\ell}{2} + 2\ell - 3 = \binom{\ell-1}{3} - \binom{\ell-2}{2}. \blacksquare$$

4.5 Proposition:

If $r > 3$ and $a = a_i - a_j$, $1 < i < j \leq \ell$. Then

1. $d_3(a_{m_1} a_i a_j) = d_3(a_i a_{m_2} a_j) = d_3(a_i a_j a_{m_3}) = O_{A_4}$, for each $1 \leq m_1 < i < m_2 < j < m_3 \leq \ell$
2. $\text{Im } d_2 = \ker d_3$.
3. $H^2(A(\mathcal{A}), a) = 0$.
4. If $r > 3$, $\dim(\text{Im } d_3) = \binom{\ell}{3} - \binom{\ell-2}{2}$.
5. If $r = 3$, $\dim(\text{Im } d_3) = \binom{\ell-2}{2}$.
6. If $r = 3$, then $\dim H^3(A(\mathcal{A}), a) = \binom{\ell-1}{3} - \binom{\ell-2}{2}$.

Proof:

To prove our claim, we need to study the homomorphism $d_3: A_3 \rightarrow A_4$, which is defined by; $d_3(a_{m_1} a_{m_2} a_{m_3}) = a_{m_1} a_{m_2} a_{m_3} a_i - a_{m_1} a_{m_2} a_{m_3} a_j$, for all 3 – NBC-monomials $a_{m_1} a_{m_2} a_{m_3} \in A_3$, since the collection $\{a_{m_1} a_{m_2} a_{m_3}; \{H_{m_1}, H_{m_2}, H_{m_3}\} \in \text{NBC}_3(\mathcal{A})$, where $1 \leq m_1 < m_2 < m_3 \leq \ell\}$ play as a basis for A_3 as a K –module. So, we will discuss the following possible cases:-

Case (1):- If $\{m_1, m_2, m_3\} = \{i, j, m\}$, then we have;

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = \begin{cases} d_3(a_m a_i a_j), & 1 \leq m < i < j \leq \ell \\ d_3(a_i a_m a_j), & 1 \leq i < m < j \leq \ell = O_{A_4}, \\ d_3(a_i a_j a_m), & 1 \leq i < j < m \leq \ell \end{cases}$$

for this case, the number of such choices is $\binom{\ell-2}{1} = \ell - 2$.

Case (2):- If m_1 or m_2 or $m_3 = i$ and $\neq j$, then;

$$\begin{aligned} d_3(a_{m_1} a_{m_2} a_{m_3}) &= \begin{cases} d_3(a_{m_1} a_{m_2} a_i); & 1 \leq m_1 < m_2 < i < j \leq \ell \\ d_3(a_{m_1} a_i a_{m_3}); & 1 \leq m_1 < i < m_3 \leq \ell \\ d_3(a_i a_{m_2} a_{m_3}); & 1 \leq i < m_2 < m_3 \leq \ell \end{cases} \\ &= \begin{cases} -a_{m_1} a_{m_2} a_i a_j; & 1 \leq m_1 < m_2 < i < j \leq \ell \\ -a_{m_1} a_i a_{m_3} a_j; & 1 \leq m_1 < i < m_3 \leq \ell \\ -a_i a_{m_2} a_{m_3} a_j; & 1 \leq i < m_2 < m_3 \leq \ell \end{cases} \\ &= \begin{cases} -a_{m_1} a_{m_2} a_i a_j; & 1 \leq m_1 < m_2 < i < j \leq \ell \\ \begin{cases} -a_{m_1} a_i a_{m_3} a_j; & 1 \leq m_1 < i < m_3 < j \leq \ell \\ a_{m_1} a_i a_j a_{m_3}; & 1 \leq m_1 < i < j < m_3 \leq \ell \end{cases} \\ \begin{cases} -a_i a_{m_2} a_{m_3} a_j; & 1 \leq i < m_2 < m_3 < j \leq \ell \\ a_i a_{m_2} a_j a_{m_3}; & 1 \leq i < m_2 < j < m_3 \leq \ell \\ -a_i a_j a_{m_2} a_{m_3}; & 1 \leq i < j < m_2 < m_3 \leq \ell \end{cases} \end{cases} \end{aligned}$$



$$= \begin{cases} -d_3(a_{m_1} a_{m_2} a_j); & 1 \leq m_1 < m_2 < i < j \leq \ell \\ \begin{cases} -d_3(a_{m_1} a_{m_3} a_j); & 1 \leq m_1 < i < m_3 < j \leq \ell \\ d_3(a_{m_1} a_j a_{m_3}); & 1 \leq m_1 < i < j < m_3 \leq \ell \end{cases} \\ \begin{cases} -d_3(a_{m_2} a_{m_3} a_j); & 1 \leq i < m_2 < m_3 < j \leq \ell \\ d_3(a_{m_2} a_j a_{m_3}); & 1 \leq i < m_2 < j < m_3 \leq \ell \end{cases} \\ -d_3(a_j a_{m_2} a_{m_3}); & 1 \leq i < j < m_2 < m_3 \leq \ell \end{cases}$$

- If $r = 4$, we will discuss all cases given in equations (2.):-

Case (2.1):- If $1 \leq m_1 < m_2 < i < j \leq \ell$, then $\{H_1, H_{m_1}, H_{m_2}, H_i, H_j\}$ is a dependent subarrangement of \mathcal{A} . Thus, $\partial_5(e_1 e_{m_1} e_{m_2} e_i e_j) \in I_4$ and;

$$\begin{aligned} d_3(a_{m_1} a_{m_2} a_i) &= -a_1 a_{m_2} a_i a_j + a_1 a_{m_1} a_i a_j - a_1 a_{m_1} a_{m_2} a_j + a_1 a_{m_1} a_{m_2} a_i \\ &= d_3(a_1 a_{m_2} a_i) - d_3(a_1 a_{m_1} a_i) + d_3(a_1 a_{m_1} a_{m_2}). \end{aligned}$$

Case (2.2):- If $1 \leq m_1 < i < m_3 < j \leq \ell$, then the subarrangement $\{H_1, H_{m_1}, H_i, H_{m_3}, H_j\}$ of \mathcal{A} is a dependent subarrangement. Thus, $\partial_5(e_1 e_{m_1} e_i e_{m_3} e_j) \in I_4$ and;

$$\begin{aligned} d_3(a_{m_1} a_i a_{m_3}) &= -a_1 a_{m_2} a_i a_j + a_1 a_{m_1} a_i a_j - a_1 a_{m_1} a_{m_2} a_j + a_1 a_{m_1} a_{m_2} a_i = \\ &= d_3(a_1 a_i a_{m_3}) + d_3(a_1 a_{m_1} a_i) - d_3(a_1 a_{m_1} a_{m_3}). \end{aligned}$$

Case (2.3):- If $1 \leq m_1 < i < j < m_3 \leq \ell$, then the subarrangement $\{H_1, H_{m_1}, H_i, H_j, H_{m_3}\}$ of \mathcal{A} is a dependent subarrangement. Thus, $\partial_5(e_1 e_{m_1} e_i e_j e_{m_3}) \in I_4$ and;

$$\begin{aligned} d_3(a_{m_1} a_i a_{m_3}) &= a_1 a_{m_2} a_i a_j - a_1 a_{m_1} a_i a_j + a_1 a_{m_1} a_{m_2} a_j - a_1 a_{m_1} a_{m_2} a_i = d_3(a_1 a_i a_{m_3}) + \\ &= d_3(a_1 a_{m_1} a_i) - d_3(a_1 a_{m_1} a_{m_3}). \end{aligned}$$

Case (2.4):- If $1 \leq i < m_2 < m_3 < j \leq \ell$, then the subarrangement $\{H_1, H_i, H_{m_2}, H_{m_3}, H_j\}$ of \mathcal{A} is a dependent subarrangement. Thus, $\partial_5(e_1 e_i e_{m_2} e_{m_3} e_j) \in I_4$ and;

$$\begin{aligned} d_3(a_{m_1} a_i a_{m_3}) &= -a_i a_{m_2} a_{m_3} a_j = -a_1 a_{m_2} a_{m_3} a_j + a_1 a_i a_{m_3} a_j - a_1 a_i a_{m_2} a_j + \\ &= a_1 a_i a_{m_2} a_{m_3} = d_3(a_1 a_i a_{m_3}) + d_3(a_1 a_i a_{m_2}) - d_3(a_1 a_{m_2} a_{m_3}). \end{aligned}$$

Case (2.5):- If $1 \leq i < m_2 < j < m_3 \leq \ell$, then the subarrangement $\{H_1, H_i, H_{m_2}, H_j, H_{m_3}\}$ of \mathcal{A} is a dependent subarrangement. Thus, $\partial_5(e_1 e_i e_{m_2} e_j e_{m_3}) \in I_4$ and;

$$\begin{aligned} d_3(a_{m_1} a_i a_{m_3}) &= a_i a_{m_2} a_j a_{m_3} = a_1 a_{m_2} a_j a_{m_3} - a_1 a_i a_j a_{m_3} + a_1 a_i a_{m_2} a_{m_3} - a_1 a_i a_{m_2} a_j = \\ &= d_3(a_1 a_i a_{m_3}) - d_3(a_1 a_i a_{m_2}) + d_3(a_1 a_{m_2} a_{m_3}). \end{aligned}$$

Case (2.6):- If $1 \leq i < j < m_2 < m_3 \leq \ell$, then the subarrangement $\{H_1, H_i, H_j, H_{m_2}, H_{m_3}\}$ of \mathcal{A} is a dependent subarrangement. Thus, $\partial_5(e_1 e_i e_j e_{m_2} e_{m_3}) \in I_4$ and;

$$\begin{aligned} d_3(a_{m_1} a_i a_{m_3}) - a_1 a_j a_j a_{m_3} + a_1 a_{m_1} a_{m_2} a_{m_3} - a_1 a_i a_j a_{m_3} + a_1 a_i a_j a_{m_2} &= d_3(a_1 a_i a_{m_3}) - \\ &= d_3(a_1 a_i a_{m_2}) + d_3(a_1 a_{m_2} a_{m_3}). \end{aligned}$$

For this case, the number of such repetition is equal to the number of our choices of 2 –indices equal i and not equal j from ℓ choices which is equal to $\binom{\ell-2}{3}$.

Case(3):- If m_1 or m_2 or $m_3 \neq i$ or j , then;

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = a_{m_1} a_{m_2} a_{m_3} a_i - a_{m_1} a_{m_2} a_{m_3} a_j$$

$$= \begin{cases} a_{m_1} a_{m_2} a_{m_3} a_i^{-1} a_{m_1} a_{m_2} a_{m_3} a_j; & 1 \leq m_1 < m_2 < m_3 < i < j \leq \ell \\ -a_{m_1} a_{m_2} a_i a_{m_3}^{-1} a_{m_1} a_{m_2} a_{m_3} a_j; & 1 \leq m_1 < m_2 < i < m_3 < j \leq \ell \\ -a_{m_1} a_{m_2} a_i a_{m_3} + a_{m_1} a_{m_2} a_j a_{m_3}; & 1 \leq m_1 < m_2 < i < j < m_3 \leq \ell \\ a_{m_1} a_i a_{m_2} a_{m_3}^{-1} a_{m_1} a_{m_2} a_{m_3} a_j; & 1 \leq m_1 < i < m_2 < m_3 < j \leq \ell \\ a_{m_1} a_i a_{m_2} a_{m_3} + a_{m_1} a_{m_2} a_j a_{m_3}; & 1 \leq m_1 < i < m_2 < j < m_3 \leq \ell \\ -a_{m_1} a_i a_{m_2} a_{m_3}^{-1} a_{m_1} a_j a_{m_2} a_{m_3}; & 1 \leq m_1 < i < j < m_2 < m_3 \leq \ell \\ -a_i a_{m_1} a_{m_2} a_{m_3}^{-1} a_{m_1} a_{m_2} a_{m_3} a_j; & 1 \leq i \leq m_1 < m_2 < m_3 < j \leq \ell \\ -a_i a_{m_1} a_{m_2} a_{m_3} + a_{m_1} a_{m_2} a_j a_{m_3}; & 1 \leq i \leq m_1 < m_2 < j < m_3 \leq \ell \\ -a_i a_{m_1} a_{m_2} a_{m_3}^{-1} a_{m_1} a_j a_{m_2} a_{m_3}; & 1 \leq i \leq m_1 < j < m_2 < m_3 \leq \ell \\ -a_i a_{m_1} a_{m_2} a_{m_3} + a_j a_{m_1} a_{m_2} a_{m_3}; & 1 \leq i < j \leq m_1 < m_2 < m_3 \leq \ell \end{cases}$$

- If $r > 4$, then every subarrangement $\{H_{n_1}, H_{n_2}, H_{n_3}, H_{n_4}\}$ with $1 \leq n_1 < n_2 < n_3 < n_4 \leq \ell$ forms a 4 – NBC basis of \mathcal{A} , hence $d_3(a_{m_1} a_{m_2} a_{m_3})$ can be written as a linear combination of NBC-monomial. So, there is no repetition among them and the number of such cases is equal to all the choices of the indices $m_1 < m_2 < m_3$ from ℓ choices such that m_1, m_2 and m_3 are not i or j , that is $\binom{\ell-2}{3}$.
- If $r = 4$, then for every cases above we will rewrite $d_3(a_{m_1} a_{m_2} a_{m_3})$ as a linear combination of NBC-monomials by using the fact that every subarrangement $\{H_1, H_{n_1}, H_{n_2}, H_{n_3}\}$ with $1 \leq n_1 < n_2 < n_3 \leq \ell$ forms 4 – NBC base for \mathcal{A} , and the fact that every $\{H_1, H_i, H_{m_1}, H_{m_2}, H_{m_3}\}$ and $\{H_1, H_j, H_{m_1}, H_{m_2}, H_{m_3}\}$ are 5 – dependent subarrangement of \mathcal{A} for all $1 \leq m_1 < m_2 < m_3 \leq \ell$ and not equal to i or j .

Notice that, if $m_1 = 1$ for the cases (3.1) to (3.6), then $d_3(a_{m_1} a_{m_2} a_{m_3})$ can be written as a linear combination of NBC-monomail, so we shall assume $m_1 > 1$ for the other cases and by a simple calculations we get:

$$d_3(a_{m_1} a_{m_2} a_{m_3}) = d_3(a_1 a_{m_2} a_{m_3}) - d_3(a_1 a_{m_1} a_{m_3}) + d_3(a_1 a_{m_1} a_{m_2}).$$

Now we will prove our claim as follows:

For (1):- It is clear from Case (1).

For (2):- From the fact that $Im d_2 \subseteq \ker d_3$ and from case (1) we have each of $a_m a_i a_j = d_2(a_m a_i)$, $a_i a_m a_j = d_2(a_i a_m)$, $a_i a_j a_m = d_2(i a_m) \in Im d_2$. So $\ker d_3 = Im d_2$.

For (3):- $H^2(A(\mathcal{A}), a) = \frac{\ker d_3}{Im d_2} = 0$.

For (4):- If $r > 4$, from our discussion in the cases 2 and 3 above, we have $Im d_3$ will be generated by all the 3 – NBC- monomials $a_{m_1} a_{m_2} a_{m_3}$ such that; There are no two choices of them can be i and j ; and by deleting the 3 – NBC- monomials that satisfied two of a_{m_1}, a_{m_2} and a_{m_3} are not a_i or a_j and by deleting the 3 – NBC-monomials that satisfied two of a_{m_1}, a_{m_2} and a_{m_3} are not a_i or a_j and the third is a_j , where such monomials represent their repetition that we discussed in case 2 above. So $\dim Im d_3 = \binom{\ell}{3} - \binom{\ell-2}{2}$, where the number of repetition is equal to $\binom{\ell}{2} - \binom{\ell-1}{2} - (\ell - 2) = \binom{\ell-2}{2}$.



For (5):- If $r = 4$, then $Im d_3$ has a basis, the set of all 3 – NBC-monomials that begin with a_1 plays as a rule to construct it by removing that containing a_j , so $\dim Im d_3 = \binom{\ell-2}{2}$.

For (6):- If $r = 4$, then $d_4: A_4 \rightarrow 0$ has $\ker d_4 = A_4$ so $\dim \ker d_4 = \binom{\ell-1}{3}$. From (5) above we have $H^3(A(\mathcal{A}), a) = \dim \ker d_4 - \dim Im d_3 = \binom{\ell-1}{3} - \binom{\ell-2}{2}$. ■

5.A GENERALIZATION OF OUR WORK:

In this section, we will generalize our work as follows:

5.1 Theorem:

If $a = a_1 - a_j, 2 \leq j \leq \ell$, then :

1. $H^{r-2}(A(\mathcal{A}), a) = 0$.
2. $\dim(Im d_{r-1}) = \binom{\ell-2}{r-2}$.
3. $\dim(H^{r-1}(A(\mathcal{A}), a)) = \binom{\ell-1}{r-1} - \binom{\ell-2}{r-2}$.

Proof:

The \mathcal{K} –homomorphism $d_{r-1}: A_{r-1} \rightarrow A_r$ is defined as;

$$d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) = a_{m_1} \cdots a_{m_{r-1}} a_1 - a_{m_1} \cdots a_{m_{r-1}} a_j ;$$

for all NBC-monomial related to $(r - 1) -$ section $\{H_{m_1}, \dots, H_{m_{r-1}}\}$ of Π . We will study all the possible cases for our choices to $m_1 < m_2 < \dots < m_{r-1} \leq \ell$, as follows:-

Case (1):- If $m_1 = 1$ and $m_k = j$ for some $2 \leq k \leq r - 1$, then;

$$d_{r-1}(a_1 \cdots a_{m_{r-1}}) = a_1 \cdots a_{m_{r-1}} a_1 - a_1 \cdots a_j \cdots a_{m_{r-1}} a_j = 0_{A_r} - 0_{A_r} = 0_{A_r}.$$

Now, we can prove (1) above as; if $\{H_{m_1}, \dots, H_{m_{r-1}}\}$, then $a_1 \cdots a_{m_{r-1}} \in \ker d_{r-1}$. We notice that $d_{r-1}(a_1 \cdots \hat{a}_j \cdots a_{m_{r-1}} a_j) = (-1)^{m+1} a_1 \cdots a_j \cdots a_{m_{r-1}}$, where m is a positive integer represents the number of transposition that we needed to rearrange $a_{m_1} \cdots a_{m_{r-1}}$ with $m_1 = 1 < m_2 < \dots < j < \dots < m_{r-1} \leq \ell$. Thus, $\ker d_{r-1} = Im d_{r-2}$ and our claim in (1) above is done, that is $H^{r-2}(A(\mathcal{A}), a) = \frac{\ker d_{r-1}}{Im d_{r-2}} = 0$.

Case (2):- If $m_1 = 1$ and $m_k \neq j$ for all $2 \leq k \leq r - 1$, then we have the following:-

$d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) = (-1)^{n+1} a_1 \cdots a_j \cdots a_{m_{r-1}} = (-1)^{n+1} (-1)^r d_{r-1}(a_{m_2} \cdots a_j \cdots a_{m_{r-1}})$, where n is a positive integer represents the number of the transpositions that we needed to reordered $a_1 \cdots a_{m_{r-1}} \cdots a_j$ to be $a_1 \cdots a_j \cdots a_{m_{r-1}}$ with $m_1 < m_2 < \dots < j < \dots < m_{r-1}$. We notice that $a_1 \cdots a_j \cdots a_{m_{r-1}}$ is an NBC-monomial.

Case (3):- If $m_1 \neq i$ and $m_k \neq j$ for all $1 \leq k \leq r - 1$, then, we need to compute the position of H_j in $H_1 \cdots H_{m_1} \cdots H_{m_{k-1}} H_j H_{m_k} \cdots H_{m_{r-1}}$. Since $\{H_1, H_{m_1}, \dots, H_j, \dots, H_{m_{r-1}}\}$ is an $(r + 1) -$ dependent section for Π , hence $\partial_{r+1}(e_1 e_{m_1} \cdots e_j \cdots e_{m_{r-1}}) \in I_r$, so;

$$a_{m_1} \cdots a_j \cdots a_{m_{r-1}} - a_1 a_{m_2} \cdots a_j \cdots a_{m_{r-1}} + a_1 a_{m_1} a_{m_3} \cdots a_j \cdots a_{m_{r-1}} + \dots + (-1)^{r+1} a_1 a_{m_2} \cdots a_j \cdots a_{m_{r-2}} = 0_{A_r}$$



we define the homomorphism $d_{r-1}: A_{r-1} \rightarrow A_r$ as follows:-

$$d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) = a_{m_1} \cdots a_{m_{r-1}} a_1 - a_{m_1} \cdots a_{m_{r-1}} a_j = (-1)^{r-1} a_1 a_{m_1} \cdots a_{m_{r-1}} + (-1)^{r-k+1} a_{m_1} \cdots a_j \cdots a_{m_{r-1}} \dots (4.1.1).$$

Thus;

$$\begin{aligned} & (-1)^0 \widehat{a_1} a_{m_1} \cdots a_j \cdots a_{m_{r-1}} + (-1)^1 a_1 \widehat{a_{m_1}} \cdots a_j \cdots a_{m_{r-1}} + \cdots + \\ & (-1)^{k-1} a_1 \cdots \widehat{a_{m_{k-1}}} a_j \cdots a_{m_{r-1}} + (-1)^k a_1 \cdots a_{m_{k-1}} \widehat{a_j} a_{m_k} \cdots a_{m_{r-1}} + \\ & (-1)^k a_1 \cdots a_{m_{k-1}} a_j \widehat{a_{m_k}} a_{m_{k+1}} \cdots a_{m_{r-1}} + \cdots + (-1)^r a_1 \cdots a_j \cdots a_{m_{r-2}} \widehat{a_{m_{r-1}}} = 0_{A_r} \dots (4.1.2) \end{aligned}$$

So, from the equation (4.1.2); we have

$$\begin{aligned} & \widehat{a_1} a_{m_1} \cdots a_j \cdots a_{m_{r-1}} = (-1)^2 a_1 \widehat{a_{m_1}} \cdots a_j \cdots a_{m_{r-1}} + \cdots + \\ & (-1)^k a_1 \cdots \widehat{a_{m_{k-1}}} a_j a_{m_k} \cdots a_{m_{r-1}} + (-1)^{k+1} a_1 \cdots a_{m_{k-1}} \widehat{a_j} a_{m_k} \cdots a_{m_{r-1}} + \cdots + \\ & (-1)^{r+1} a_1 \cdots a_j \cdots a_{m_{r-2}} \widehat{a_{m_{r-1}}} \dots (4.1.3). \end{aligned}$$

By substituting the equation (4.1.3) in the equation (4.1.2), we have;

$$\begin{aligned} d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) &= (-1)^{r-1} a_1 \cdots a_{m_{k-1}} \widehat{a_j} a_{m_k} \cdots a_{m_{r-1}} + \\ & (-1)^{r-k+3} a_1 \widehat{a_{m_1}} a_{m_2} \cdots a_j \cdots a_{m_{r-1}} + (-1)^{r-k+4} a_1 a_{m_1} \widehat{a_{m_2}} \cdots a_j \cdots a_{m_{r-1}} + \cdots + \\ & (-1)^{r-k+1+k+1} a_1 \cdots a_{m_{k-1}} \widehat{a_j} a_{m_k} \cdots a_{m_{r-1}} + \cdots + (-1)^{r-k-1+r+1} a_1 \cdots a_j \cdots a_{m_{r-2}} \widehat{a_{m_{r-1}}}. \end{aligned}$$

$$\begin{aligned} d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) &= (-1)^{r-k+1} d_{r-1}(a_1 \widehat{a_{m_1}} a_{m_2} \cdots a_{m_{r-1}}) + \\ & (-1)^{r-k+4} d_{r-1}(a_1 a_{m_1} \widehat{a_{m_2}} a_{m_3} \cdots a_{m_{r-1}}) + \cdots + (-1)^{2r-k} d_{r-1}(a_1 a_{m_1} \cdots a_{m_{r-2}} \widehat{a_{m_{r-1}}}). \end{aligned}$$

Now to prove our conjuctor:

For (2):- That is, $Im d_{r-1}$ will be generated by the $(r - 1) - NBC$ -monomials that begin with a_1 and contains no a_j and the number of such monomials is $\binom{\ell-2}{r-2}$, the number of all $(r - 2) - NBC$ -monomials that not contain each of a_1 and a_j , that is; $\dim Im d_{r-1} = \binom{\ell-2}{r-2}$.

For (3):- Since $d_r: A_r \rightarrow 0$ is the zero homomorphism, hence $\ker d_r = A_r$ and $\dim \ker d_r = \binom{\ell-1}{r-1}$. Therefore, $\dim H^{r-1}(A(\mathcal{A}), a) = \dim \ker d_r - \dim Im d_{r-1} = \binom{\ell-1}{r-1} - \binom{\ell-2}{r-2}$.

5.2 Theorem:

If $a = a_1 - a_j, 2 < j \leq \ell$, and for $3 \leq k < r - 1$, we have;

1. $H^{k-1}(A(\mathcal{A}), a) = 0$.
2. $\dim(Im d_k) = \binom{\ell}{k} - \binom{\ell-2}{k-1}$.

Proof:-

We will use the induction to prove our claim and we will recall proposition (3.3) to show that our conjecture is true for $k = 3$. Assume our claim is true for $k - 1$ and we will prove it for k . So, we shall begin with the definition of $d_k: A_k \rightarrow A_{k+1}$. Since the set of all $K - NBC$ -monomials play as a basis of $A_k(\mathcal{A})$, so firstly we will use our ordering to arrange the position of H_j in the HP Π . To explain it's position in any NBC -base contain it, let $\{H_{m_1}, \dots, H_{m_k}\}$ be an NBC base such that $H_{m_n} \neq H_j$ for all $1 \leq n \leq k$, then we can assume

the position of H_j among H_{m_1}, \dots, H_{m_k} that respect the hypersolvable ordering that we defined on the hyperplanes of \mathcal{A} is;

$$H_{m_1}, \dots, H_{m_n}, H_j, H_{m_{n+1}}, \dots, H_{m_k},$$

such that;

$$\underbrace{m_1 < m_2 < \dots < m_n}_{n+1} < j < \underbrace{m_{n+1} < \dots < m_k}_{k-n}$$

$$\text{Thus, } d_k(a_{m_1} \dots a_{m_k}) = a_{m_1} \dots a_{m_k} a_1 - a_{m_1} \dots a_{m_k} a_j;$$

and we will study the possible cases of our choices of m_1, \dots, m_k , as follows:-

Case (1):- If $m_1 = 1$ and $m_n = j$ for some $1 \leq n \leq k$, then;

$$d_k(a_1 a_2 \dots a_{m_k}) = a_1 \dots a_{m_k} a_1 - a_1 \dots a_2 \dots a_{m_k} a_j = 0_{A_{k+1}} - 0_{A_{k+1}} = 0_{A_{k+1}}.$$

Now, we can prove (1) above as:- If $H_1, H_j \in \{H_{m_1}, \dots, H_{m_k}\}$, then $a_1 \dots a_{m_k} \in \ker d_k$. We notice that $d_{k-1}(a_1 \dots \hat{a}_j \dots a_{m_k}) = -a_1 \dots \hat{a}_j \dots a_{m_k} a_j = (-1)^{m+1} a_1 \dots a_j \dots a_{m_k}$. Thus $\ker d_k = \text{Im } d_{k-1}$ and our claim in (1) is done. That is, $H^{k-1}(A(\mathcal{A}), a) = \frac{\ker d_k}{\text{Im } d_{k-1}} = 0$.

Case (2):- If $m_1 = 1$ and $m_n \neq j$ for all $2 \leq n \leq k$, then we have the following:-

$$d_k(a_1 \dots a_{m_k}) = (-1)^{n+1} a_1 \dots a_j \dots a_{m_k} = (-1)^{n+1} (-1)^k d_k(a_{m_2} \dots a_j \dots a_{m_k}),$$

where n is the positive integer represent the number of the transposition that we needed to reordered $a_1 \dots a_{m_k} \dots a_j$ to be $a_1 \dots a_j \dots a_{m_k}$ with $m_1 < m_2 < \dots < j < \dots < m_k$, we notice that $a_1 \dots a_j \dots a_{m_k}$ is an *NBC*-monomial.

Case (3):- If $m_1 \neq 1$ and $m_n \neq j$ for all $1 \leq n \leq k$, then we have the following:-

Let $\{H_{m_1}, \dots, H_{m_k}\}$ be an *NBC* base such that $H_{m_n} \neq H_j$ for all $1 \leq n \leq k$, then we can assume the position of H_j among H_{m_1}, \dots, H_{m_k} that respect the hypersolvable ordering that we defined on the hyperplanes of \mathcal{A} . Thus;

$$d_k(a_{m_1} \dots a_{m_k}) = a_{m_1} \dots a_{m_k} a_1 - a_{m_1} \dots a_{m_k} a_j = (-1)^k a_1 a_{m_1} \dots a_{m_k} + (-1)^{n-k+1} a_{m_1} \dots a_{m_n} a_j a_{m_{n+1}} \dots a_{m_k}.$$

So, $d_k(a_{m_1} \dots a_{m_k})$ can be written as a linear combination of $(k + 1) - \text{NBC}$ -bases.

For (2):- The $k - \text{NBC}$ -monomial plays as a basis to $\text{Im } d_k$ with keeping in mind that we need to remove the $k - \text{NBC}$ -monomials contain a_j . That is $\dim \text{Im } d_k = \binom{\ell}{k} - \binom{\ell-2}{k-1}$. ■

5.3 Theorem:

If $a = a_i - a_j$, where $1 < i < j \leq \ell$, then;

1. $H^{r-2}(A(\mathcal{A}), a) = 0$.
2. $\dim(\text{Im } d_r) = \binom{\ell-2}{r-2}$.
3. $\dim(H^{r-1}(A(\mathcal{A}), a)) = \binom{\ell-1}{r-1} - \binom{\ell-2}{r-2}$.

Proof:-



To show our claim, we need to study the homomorphism $d_{r-1}: A_{r-1} \rightarrow A_r$. As we know, the family of all $(r - 1) - NBC$ basis play a role to construct a basis to A_{r-1} . Then, for all $(r - 1) - NBC$ -monomial $a_{m_1} \cdots a_{m_{r-1}}$, we define d_{r-1} as;

$$d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) = a_{m_1} \cdots a_{m_{r-1}} a_i - a_{m_1} \cdots a_{m_{r-1}} a_j.$$

we will study the possible cases for our choices to the indices $1 \leq m_1 < \cdots < m_{r-1} \leq \ell$ as follows:-

Case (1): If $m_{k_1} = i$ and $m_{k_2} = j$ for some $1 \leq k_1 < k_2 \leq r - 1$, then;

$$d_{r-1}(a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2+1}} a_{m_{k_2+1}} \cdots a_{m_{r-1}}) = a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{r-1}} a_i - a_{m_1} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2+1}} \cdots a_{m_{r-1}} a_j = 0_{A_r} - 0_{A_r} = 0_{A_r}.$$

For (1):-

Since $Im d_{r-1} \subseteq \ker d_r$ and from the fact that;

$$\begin{aligned} d_{r-2}(a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} \widehat{a}_j a_{m_{k_2+1}} \cdots a_{m_{r-1}}) \\ = (-1) a_{m_1} \cdots a_i \cdots a_{m_{k_2-1}} a_{m_{k_2+1}} \cdots a_{m_{r-1}} a_j \\ = (-1)^{r-m_{k_2}} a_{m_1} \cdots a_i \cdots a_{m_{k_2-1}} a_j a_{m_{k_2+1}} a_{m_{r-1}}. \end{aligned}$$

Therefore, $a_{m_1} \cdots a_i \cdots a_j \cdots a_{m_{r-1}} \in Im d_{r-2}$ and $Im d_{r-2} = \ker d_{r-1}$. So;

$$H^{r-2}(A(\mathcal{A}), a) = \ker d_{r-1} / Im d_{r-2} = \frac{Z_{r-2}}{B_{r-2}} = 0.$$

Case (2):- If $m_{k_1} = i$ for some $1 \leq k_1 \leq r - 1$ and there is no $m_k = j$ for all $k_1 \leq k \leq r - 1$, then;

$$d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) = -a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{r-1}} a_j = (-1)^{r-k_2+1} \underbrace{a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}}}_{k_2-1} \underbrace{a_j a_{m_{k_2+1}} \cdots a_{m_{r-1}}}_{r-k_2} \dots \quad (4.3.1).$$

where $k_2 = \text{Min}\{k | H_j \trianglelefteq H_{m_k}\}$. Thus;

$$\begin{aligned} d_{r-1}(a_{m_1} \cdots a_{m_{r-1}}) \\ = (-1)^{r-k_2+1} (-1)^{r-k_1+1} d_{r-1}(a_{m_1} \cdots a_{m_{k_1-1}} \widehat{a}_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2}} \cdots a_{m_{r-1}}) \\ = (-1)^{2r-k_1-k_2} d_{r-1}(a_{m_1} \cdots a_{m_{k_1-1}} a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2}} \cdots a_{m_{r-1}}). \end{aligned}$$

As we know, $a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2}} \cdots a_{m_{r-1}}$ is not $r - NBC$ -monomial, so we need to write it as a linear combination of $r - NBC$ -monomials. In fact;

$$\{H_1, H_{m_1}, \dots, H_{m_{k_1-1}}, H_i, H_{m_{k_1+1}}, \dots, H_{m_{k_2-1}}, H_j, H_{m_{k_2}}, \dots, H_{m_{r-1}}\};$$

is a dependent subarrangement of \mathcal{A} and;

$$\begin{aligned} \partial_{r+1}(e_1 e_{m_1} \cdots e_{m_{k_1-1}} e_i e_{m_{k_1+1}} \cdots e_{m_{k_2-1}} e_j e_{m_{k_2}} \cdots e_{m_{r-1}}) \in I_r. \text{ Thus;} \\ (-1)^0 \widehat{a}_1 a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2}} \cdots a_{m_{r-1}} + \\ (-1)^1 a_1 \widehat{a}_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2}} \cdots a_{m_{r-1}} + \cdots + \\ (-1)^{k_1} a_1 a_{m_1} \cdots a_{m_{k_1-1}} \widehat{a}_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} a_j a_{m_{k_2}} \cdots a_{m_{r-1}} + \cdots + \\ (-1)^{k_2} a_1 a_{m_1} \cdots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \cdots a_{m_{k_2-1}} \widehat{a}_j a_{m_{k_2}} \cdots a_{m_{r-1}} + \end{aligned}$$



$$\begin{aligned}
 & (-1)^{k_2+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j \widehat{a_{m_{k_2}}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^r a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots \widehat{a_{m_{r-1}}} = \\
 & a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \\
 & (-1)^2 a_1 \widehat{a_{m_1}} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{k_1+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} \widehat{a_i} a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{k_2+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} \widehat{a_j} a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{r+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots \widehat{a_{m_{r-1}}} \dots (4.3.2).
 \end{aligned}$$

By substituting (4.3.2) in (4.3.1) we have:-

$$\begin{aligned}
 d_{r-1}(a_{m_1} \dots a_{m_{r-1}}) &= (-1)^{r-k_2+3} (-1)^{r-k_2} d_{r-1}(a_1 \widehat{a_{m_1}} a_{m_2} \dots a_{m_{r-1}}) + \\
 & (-1)^{r-k_2+4} (-1)^{r-k_2} d_{r-1}(a_1 a_{m_2} \widehat{a_{m_2}} a_{m_3} \dots a_{m_{r-1}}) + \dots + \\
 & (-1)^{r-k_2+k_1+2} (-1)^{r-k_2} d_{r-1}(a_1 a_{m_2} \dots a_{m_{k_1-1}} \widehat{a_i} a_{m_{k_1+1}} \dots a_{m_{r-1}} a_j) + \dots + \\
 & (-1)^{2r-k_2+2} (-1)^{r-k_2} d_{r-1}(a_1 a_{m_2} \dots a_{m_{r-2}} \widehat{a_{m_{r-1}}} a_j) = (-1) d_{r-1}(a_1 \widehat{a_{m_1}} \dots a_{m_{r-1}}) + \\
 & (-1)^2 d_{r-1}(a_1 a_{m_2} \widehat{a_{m_2}} \dots a_{m_{r-1}}) + \dots + (-1)^{k_1-1} d_{r-1}(a_1 a_{m_1} \dots \widehat{a_{k_1-1}} a_i \dots a_{m_{r-1}}) + \\
 & (-1)^{k_1+1} d_{r-1}(a_1 a_{m_1} \dots a_i \widehat{a_{k_1+1}} \dots a_{m_{r-1}}) + \dots + (-1)^{r+2} d_{r-1}(a_1 a_{m_1} \dots a_{m_{r-2}} \widehat{a_{m_{r-1}}}) + \\
 & (-1)^{k_1} a_1 a_{m_1} \dots \widehat{a_i} \dots a_{m_{r-1}} a_j + (-1)^{k_1+1} a_1 a_{m_1} \dots \widehat{a_j} \dots a_{m_{r-1}} a_i = \\
 & (-1) d_{r-1}(a_1 \widehat{a_{m_1}} \dots a_{m_{r-1}}) + (-1)^2 d_{r-1}(a_1 a_{m_2} \widehat{a_{m_2}} \dots a_{m_{r-1}}) + \dots + \\
 & (-1)^{r+2} d_{r-1}(a_1 a_{m_1} \dots a_{m_{r-2}} \widehat{a_{m_{r-1}}}) - d_{r-1}(a_1 a_{m_1} \dots \widehat{a_i} \dots \widehat{a_j} \dots a_{m_{r-1}}).
 \end{aligned}$$

Thus, $d_{r-1}(a_{m_1} \dots a_{m_{r-1}})$ can be written as a linear combination of $r - NBC$ -monomials that related to the image of the homomorphism d_{r-1} of $(r - 1) - NBC$ -monomials that begin with a_1 . Such NBC -monomials depend on the NBC -basis that not begin with H_1 . As well as $m_k \neq j$ or $m_k \neq i$ and the number of such repetition is $\binom{\ell-2}{r-2}$.

Case (3):- If there is no $1 \leq n_1 < n_2 \leq r - 1$ such that; $a_{m_{n_1}} = a_i$ and $a_{m_{n_2}} = a_j$. Then, we will assume; $k_1 = \text{Min}\{k|H_i \subseteq H_{m_{k_1}}\}$ and $k_2 = \text{Min}\{k|H_j \subseteq H_{m_{k_2}}\}$, to define;

$$\begin{aligned}
 d_{r-1}(a_{m_1} \dots a_{m_{r-1}}) &= (-1)^{r-k_1+1} a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \\
 & (-1)^{r-k_2+1} a_{m_1} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} \dots (4.3.3);
 \end{aligned}$$

in order to write $d_{r-1}(a_{m_1} \dots a_{m_{r-1}})$ as a linear combination $r - NBC$ -monomials. Consequently, we recall the fact that each of; $\{H_1, H_{m_1}, \dots, H_{m_{k_1-1}}, H_i, H_{m_{k_1}}, \dots, H_{m_{r-1}}\}$, and $\{H_1, H_{m_1}, \dots, H_{m_{k_2-1}}, H_j, H_{m_{k_2}}, \dots, H_{m_{r-1}}\}$; are dependent subarrangements of \mathcal{A} , hence;

$$\begin{aligned}
 & \partial_{r+1}(e_1 e_{m_1} \dots e_{m_{k_1-1}} e_i e_{m_{k_1}} \dots e_{m_{r-1}}), \partial_{r+1}(e_1 e_{m_1} \dots e_{m_{k_2-1}} e_j e_{m_{k_2}} \dots e_{m_{r-1}}), \\
 & \partial_{r+1}(e_1 e_{m_1} \dots e_{m_{k_2-1}} e_j e_{m_{k_2}} \dots e_{m_{r-1}}) \in I_r.
 \end{aligned}$$

Thus;

$$\begin{aligned}
 & (-1)^0 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + (-1)^1 a_1 \widehat{a_{m_1}} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \\
 & (-1)^2 a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \dots + (-1)^{k_1} a_1 a_{m_1} \dots a_{m_{k_1-1}} \widehat{a_i} a_{m_{k_1}} \dots a_{m_{r-1}} + \\
 & (-1)^{k_1+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i \widehat{a_{m_{k_1}}} \dots a_{m_{r-1}} + \dots + (-1)^r a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots \widehat{a_{m_{r-1}}}.
 \end{aligned}$$

$$\begin{aligned}
 & a_{m_1} a_{m_2} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} = (-1)^2 a_1 \widehat{a_{m_1}} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \\
 & (-1)^3 a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{k_1+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} \widehat{a_i} a_{m_{k_1}} \dots a_{m_{r-1}} + \dots + (-1)^{r+1} a_1 a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots \widehat{a_{m_{r-1}}} \dots
 \end{aligned}$$

(4.3.4). Therefore;

$$\begin{aligned}
 & (-1)^0 a_{m_1} a_{m_2} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + (-1)^1 a_1 \widehat{a_{m_1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \\
 & (-1)^2 a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + (-1)^{k_2} a_1 a_{m_1} \dots a_{m_{k_2-1}} \widehat{a_j} a_{m_{k_2}} \dots a_{m_{r-1}} + \\
 & (-1)^{k_2+1} a_1 a_{m_1} \dots a_{m_{k_2-1}} a_j \widehat{a_{m_{k_2}}} \dots a_{m_{r-1}} + \dots + (-1)^r a_1 a_{m_1} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots \widehat{a_{m_{r-1}}}. \\
 & a_{m_1} a_{m_2} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} = (-1)^2 a_1 \widehat{a_{m_1}} a_{m_2} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{k_2} a_1 a_{m_1} \dots a_{m_{k_2-1}} \widehat{a_j} a_{m_{k_2}} \dots a_{m_{r-1}} + (-1)^{k_2+1} a_1 a_{m_1} \dots a_{m_{k_2-1}} \widehat{a_{m_{k_2}}} a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{r+1} a_1 a_{m_1} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots \widehat{a_{m_{r-1}}} \dots (4.3.5).
 \end{aligned}$$

By substituting (4.3.4) and (4.3.5) in (4.3.3) it follows that;

$$\begin{aligned}
 d_{r-1}(a_{m_1} \dots a_{m_{r-1}}) &= (-1)^{r-k_1+3} a_1 \widehat{a_{m_1}} a_{m_2} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \\
 & (-1)^{r-k_1+4} a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{r-k_1+1+k_1+1} a_1 a_{m_1} a_{m_2} \dots a_{m_{k_1-1}} \widehat{a_i} a_{m_{k_1}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{2r-k_1+2} a_1 a_{m_1} a_{m_2} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots \widehat{a_{m_{r-1}}} + \\
 & (-1)^{r-k_2+2} a_1 \widehat{a_{m_1}} a_{m_2} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \\
 & (-1)^{r-k_2+3} a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{r-k_2+k_2+1} a_1 a_{m_1} a_{m_2} \dots a_{m_{k_2-1}} \widehat{a_j} a_{m_{k_2}} \dots a_{m_{r-1}} + \dots + \\
 & (-1)^{r-k_2+r+1} a_1 a_{m_1} a_{m_2} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots \widehat{a_{m_{r-1}}} = (-1)^{r+3} a_1 \widehat{a_{m_1}} a_{m_2} \dots a_{m_{r-1}} a_i + \\
 & (-1)^{r+2} a_1 \widehat{a_{m_1}} a_{m_2} \dots a_{m_{r-1}} a_j + (-1)^{r+4} a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{r-1}} a_i + \\
 & (-1)^{r+3} a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{r-1}} a_j + \dots + (-1)^{2r+2} a_1 a_{m_1} a_{m_2} \dots \widehat{a_{m_{r-1}}} a_i + \\
 & (-1)^{r+1} a_1 a_{m_1} a_{m_2} \dots \widehat{a_{m_{r-1}}} a_j = (-1)^r d_{r-1}(a_1 \widehat{a_{m_1}} \dots a_{m_{r-1}}) + \\
 & (-1)^r d_{r-1}(a_1 a_{m_1} \widehat{a_{m_2}} \dots a_{m_{r-1}}) + \dots + (-1)^{2r} d_{r-1}(a_1 a_{m_1} \dots \widehat{a_{m_{r-1}}}).
 \end{aligned}$$

Therefore, every coboundary of $A_r(\mathcal{A})$ can be written as a linear combination of $r - \text{coboundaries}$ that begin with a_1 . However, the number of such NBC -monomials depends of $(r - 2)$ - NBC -monomials that begin with a_1 and contains no a_j .

For (2):- From above we established $Im d_{r-1}$ structure with;

$$\dim Im d_{r-1} = \dim B_{r-1} = \binom{\ell-2}{r-2}.$$

For (3):- As well as; $H^{r-1}(A(\mathcal{A}), a) = \frac{\ker d_r}{Im d_{r-1}} = Z_{r-1}/B_{r-1}$ and;

$$\dim H^{r-1}(A(\mathcal{A}), a) = \dim Z_{r-1} - \dim B_{r-1} = \binom{\ell-1}{r-1} - \binom{\ell-2}{r-2}. \blacksquare$$

5.4 Theorem:

If $a = a_i - a_j$, where $1 < i < j \leq \ell$, then, for $3 \leq k < r - 1$, we have:-

1. $H^{k-1}(A(\mathcal{A}), a) = 0$.
2. $\dim(Im d_k) = \binom{\ell}{k} - \binom{\ell-2}{k-1}$.

Proof:

We will prove our conjecture inductively. Recall the proposition (3.5) for $k = 3$. So, we shall assume our claim is true for $k - 1$ and we will prove it for k . In fact, the family of all $k - NBC$ -bases of \mathcal{A} plays a role to construct a basis to $A_k(\mathcal{A})$. Hence we will study the homomorphism $d_k: A_k \rightarrow A_{k+1}$ as follows; for all $k - NBC$ -monomial a_{m_1}, \dots, a_{m_k} we have;

$$d_k(a_{m_1} \dots a_{m_k}) = a_{m_1} \dots a_{m_k} a_i - a_{m_1} \dots a_{m_k} a_j.$$

The image of d_k depends of the $k - NBC$ -monomial. So, we will study all the possible cases of these NBC -monomials.

Case(1):- If there is $1 < k_1 < k_2 < r - 1$ such that $m_{k_1} = i$ and $m_{k_2} = j$, then;

$$d_k(a_{m_1} \dots a_{m_k}) = a_{m_1} \dots a_i \dots a_j \dots a_{m_k} a_i - a_{m_1} \dots a_i \dots a_j \dots a_{m_k} a_j = 0_{A_{k+1}} - 0_{A_{k+1}} = 0_{A_{k+1}}.$$

Inductively, we show that such $k - NBC$ -monomial is $k - cocycles$. So, $a_{m_1} \dots a_i \dots a_j \dots a_{m_k} \in \ker d_k$. Moreover, $Im d_{k-1} \subseteq \ker d_k$ and $a_{m_1} \dots a_i \dots a_j \dots a_{m_k} \in Im d_{k-1} = B_{k-1}$, since;

$$d_{k-1}(a_{m_1} \dots a_i \dots \hat{a}_j \dots a_{m_k}) = (-1)^{k_1-k_2} a_{m_1} \dots a_i \dots a_j \dots a_{m_k}.$$

Therefore, $\ker d_k = Im d_{k-1} (Z_{k-1} = B_{k-1})$ and $H^{k-1}(A(\mathcal{A}), a) = 0$. That is (1) above is verified.

Case (2):- If there is $1 < k_1 \leq k$ such that $m_{k_1} = i$ and there is no $1 < n \leq k$ such that $m_n = j$; then, put $k_2 = Min\{n | H_i \cong H_{m_n}\}$. Thus;

$$\begin{aligned} d_k(a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_k}) &= \\ (-1)^{k-k_2} a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_k} &= \\ (-1)^{k-k_2} (-1)^{k-k_1+2} d_k(a_{m_1} \dots a_{m_{k_1-1}} \hat{a}_i a_{m_{k_1+1}} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_k}). \end{aligned}$$

The important point here, $d_k(a_{m_1} \dots a_i \dots a_{m_k})$ is a $(k + 1) - NBC$ -monomial.

Case (3):- If there is no $1 \leq n_1 < n_2 \leq k$ such that $n_1 = i$ and $n_2 = j$, then put $k_1 = Min\{n | H_i \cong H_{m_n}\}$, and $k_2 = Min\{n | H_j \cong H_{m_n}\}$. Thus;

$$\begin{aligned} d_k(a_{m_1} \dots a_{m_k}) &= (-1)^{k-k_1+1} a_{m_1} \dots a_{m_{k_1-1}} a_i a_{m_{k_1}} \dots a_{m_k} + \\ (-1)^{k-k_2+2} a_{m_1} \dots a_{m_{k_2-1}} a_j a_{m_{k_2}} \dots a_{m_k}. \end{aligned}$$

That is, $d_k(a_{m_1} \dots a_{m_k})$ written as a linear combination of $(k + 1) - NBC$ -monomials.

For (2): Our study to the three cases above, leads to $Im d_k$ have a basis generated by $k - NBC$ -monomials that represent repetition in case (2) and co-cycle in case (1). Therefore, $\dim(Im d_k) = \binom{\ell}{k} - \binom{\ell-2}{k-1}$. ■

References

- [1] S. Roberts, On the Figures formed by the Intercepts of a System of Straight Lines in a, Plane, and on analogous relations in Space of Three Dimensions, *Proceedings of the London Mathematical Society* s1-19 (1887) 405-422.



- [2] P. Orlik and H. Terao, Arrangements of Hyperplanes, Springer Berlin Heidelberg, 1992.
- [3] E. Fadell and L. Neuwirth, Configuration Spaces., *MATHEMATICA SCANDINAVICA* 10 (1962) 111.
- [4] R. Fox and L. Neuwirth, The Braid Groups., *MATHEMATICA SCANDINAVICA* 10 (1962) 119.
- [5] E. Brieskorn, "Sur les groupes de tresses [daprès V. I. Arnold]," in *Lecture Notes in Mathematics*, Springer Berlin Heidelberg, 1973, 21-44. .
- [6] P. Orlik and L. Solomon, Combinatorics and topology of complements of hyperplanes, *Inventiones Mathematicae* 56 (1980) 167-189.
- [7] A. Hattori, Topology of C^n minus a finite number of affine hyperplanes in general position, *Journal of the Faculty of Science, the University of Tokyo. Sect. I A, Mathematics* 22 (1975) 205-219.
- [8] M. Jambu and S. Papadima, A generalization of fiber-type arrangements and a new deformation method, *Topology* 37 (1998) 1135-1164.
- [9] M. Jambu and Ş. Papadima, Deformations of hypersolvable arrangements, *Topology and its Applications* 118 (2002) 103-111.
- [10] R. P. Stanley, Supersolvable lattices, *Algebra Universalis* 2 (1972) 197-217.
- [11] H. Ali, A Topological Interpretation for Vanishing of Higher Homotopy Group of a Hypersolvable Arrangement, *t, J. Basrah Researches (Sci)*, 5 (2006) 414-433.
- [12] A. Al-Ta'ai and H. Ali, The Broken Circuit Complex and Hypersolvable Partition Complex., *Thi-Qar Sci* 2 (2010)
- [13] A. H. AL-Ta'a, H. M. Ali and M. A. Majeed, On the Orlik-Solomon algebra of a hypersolvable arrangement, *Basrah Researches (Sci)* 36 (2009) 414-433.
- [14] T. Zaslavsky, Counting the faces of cut-up spaces, *Bulletin of the American Mathematical Society* 81 (1975) 916-919.
- [15] S. Yuzvinsky, Cohomology of the Orlik-Solomon Algebras, *communication in algebras* 23 (1995) 5339-5354.
- [16] K. J. Pearson, Cohomology of OS algebras for quadratic arrangements, *Lecturas Matemáticas* 22 (2001) 103-134.

دراسة لكهمولوجيا الأورلاك-سولومون الجبراً كموديول حر لترتيبة عامة

نجوان عدنان منصور هناء مرتضى علي
جامعة البصرة / كلية العلوم / قسم الرياضيات



المستخلص

كُرسَ هذا البحث لدراسة كهمولوجيا الأورلاك-سولومون الجبر $A(\mathcal{A})$ كموديول حر لترتيبة ℓ - عامة، حيث $a = a_i - a_j$ ، $1 \leq i < j \leq \ell$. على وجه الخصوص. كما بعد $H^k(A(\mathcal{A}), a)$ تم حسابه لكل $1 \leq k \leq r - 1$.

